

Periodic orbits of radially symmetric systems with a singularity: the repulsive case

Alessandro Fonda and Rodica Toader

ABSTRACT. We study radially symmetric systems with a singularity of repulsive type. In the presence of a radially symmetric periodic forcing, we show the existence of three distinct families of subharmonic solutions: One oscillates radially, one rotates around the origin with small angular momentum, and the third one with both large angular momentum and large amplitude. The proofs are carried out by the use of topological degree theory.

1 Introduction

We are interested in proving the existence and multiplicity of periodic solutions for systems of the type

$$\ddot{x} = f(t, |x|) x. \quad (1)$$

Here, $f : \mathbb{R} \times]0, +\infty[\rightarrow \mathbb{R}$ is L^1 -Carathéodory and T -periodic with respect to its first variable, so that 0 is possibly a singularity point. We look for solutions $x(t) \in \mathbb{R}^N$ which never attain the singularity, in the sense that

$$x(t) \neq 0, \quad \text{for every } t \in \mathbb{R}. \quad (2)$$

Since system (1) is radially symmetric, the orbit of a solution always lies on a plane, so we will assume, without loss of generality, that $N = 2$.

Let us consider, e.g., the model equation

$$\ddot{x} + c(t) \frac{x}{|x|^\gamma} = e(t) \frac{x}{|x|}. \quad (3)$$

Here, $\gamma > 1$, and $c, e \in L^1_{loc}(\mathbb{R})$ are T -periodic. Notice that, if c and e are constant functions, with $e(t)$ identically zero, and $\gamma = 3$, then (3) is the equation modelling the motion of an electrical charge in the field of another charge standing still at the origin. According to the sign of c , we can have both the cases of attractive and repulsive forces. In [8] we have treated the attractive case. In this paper, we are interested in the case of a repulsive force.

Writing the solutions in polar coordinates, we see that equation (3) is equivalent to the system

$$\begin{cases} \ddot{\rho} - \frac{\mu^2}{\rho^3} + \frac{c(t)}{\rho^{\gamma-1}} = e(t), \\ \rho^2 \dot{\varphi} = \mu, \end{cases} \quad (4)$$

where μ is the (scalar) angular momentum. Let us denote by \bar{e} the mean value of the forcing term:

$$\bar{e} = \frac{1}{T} \int_0^T e(t) dt.$$

If $c(t)$ is strictly negative, integrating the first equation in (4), it is easily seen that a necessary condition for the existence of periodic solutions is that \bar{e} be negative. On the other hand, when $\gamma \geq 2$ and $\mu = 0$, it was proved in [11, 10, 5, 7, 1] that, if \bar{e} is negative, then system (4) has periodic radial solutions. Our interest is to find out whether, besides these radial solutions, there are periodic solutions rotating around the origin, as well.

In order to have an insight on the possible behaviour of the solutions, let us now give a description of the circular orbits of (3) in the case when $c(t) = \bar{c}$ and $e(t) = \bar{e}$ are negative constants. We have to look for a constant ρ satisfying

$$-\frac{\mu^2}{\rho^3} + \frac{\bar{c}}{\rho^{\gamma-1}} = \bar{e}. \quad (5)$$

Defining

$$\hat{\rho} := \left(\frac{\bar{c}}{\bar{e}} \right)^{\frac{1}{\gamma-1}},$$

it is easily seen that, for every μ , if ρ satisfies (5), then $\rho \geq \hat{\rho}$, with equality only when $\mu = 0$. Hence, there is a circle of stationary points, centered at the origin, with radius $\hat{\rho}$, and there are no circular orbits inside that circle. The circular solutions with radius ρ greater than $\hat{\rho}$ have angular momentum

$$\mu(\rho) = \sqrt{\bar{c}\rho^{4-\gamma} - \bar{e}\rho^3}.$$

Hence, the angular momentum is strictly increasing with the radius of the solutions, and

$$\lim_{\rho \rightarrow \hat{\rho}^+} \mu(\rho) = 0, \quad \lim_{\rho \rightarrow +\infty} \mu(\rho) = +\infty.$$

In particular, the circular solutions can be parametrized by the angular momentum μ , so that, in the plane (μ, ρ) , we have a curve whose projection on the first component is the whole half-line $]0, +\infty[$.

As for the period of the solutions, we have

$$\tau(\rho) = 2\pi \sqrt{\frac{\rho}{-\bar{e} + \bar{c}\rho^{1-\gamma}}},$$

so that

$$\lim_{\rho \rightarrow \hat{\rho}^+} \tau(\rho) = +\infty, \quad \lim_{\rho \rightarrow +\infty} \tau(\rho) = +\infty.$$

The function $\tau(\rho)$ has a minimum point at $\rho = \rho_{min} := \left(\frac{\bar{c}\gamma}{\bar{e}}\right)^{\frac{1}{\gamma-1}}$, with minimum value

$$\tau_{min} = \frac{2\pi}{\sqrt{\gamma-1}} \left(\frac{-\bar{c}\gamma^\gamma}{(-\bar{e})^\gamma} \right)^{\frac{1}{2(\gamma-1)}}.$$

Moreover, $\tau(\rho)$ is strictly decreasing when $\rho < \rho_{min}$ and strictly increasing when $\rho > \rho_{min}$. Hence, for every τ greater than τ_{min} , there are two circular orbits with minimal period τ : one with a small, and the other with a large angular momentum.

In the case when $c(t)$ and $e(t)$ are not necessarily constant, but T -periodic, it is natural to investigate whether (3) still has periodic solutions, rotating around the origin, having a similar behaviour to the circular orbits described above. The aim of this paper is to prove that such solutions exist, even for much more general systems like (1). The idea is to look for periodic solutions $x(t)$ whose minimal periods are sufficiently large multiples of T , and whose angular momenta are sufficiently small, or sufficiently large, respectively. In order to prevent too large eccentricities of the orbits, we will impose the radial components of the solutions $x(t)$ to be T -periodic.

Let us consider the more general system (1). It is convenient to write it in the following form:

$$\ddot{x} = \left(-h(t, |x|) + e(t) \right) \frac{x}{|x|}. \quad (6)$$

We will prove the following two theorems. The first one deals with periodic solutions having a small angular momentum.

Theorem 1 *Let the following three assumptions hold.*

(H1) *There is a function $\alpha \in L^1(0, T)$ such that*

$$\limsup_{r \rightarrow +\infty} \frac{h(t, r)}{r} \leq \alpha(t),$$

uniformly for almost every $t \in]0, T[$, and

$$\alpha(t) \leq \left(\frac{\pi}{T} \right)^2, \quad (7)$$

for almost every $t \in]0, T[$, with strict inequality on a subset of $]0, T[$ having positive measure.

(H2) *There exists a function $\eta \in L^1(0, T)$, with positive values, such that*

$$\begin{aligned} h(t, r) &\leq \eta(t), & \text{for every } r \in]0, 1] \text{ and a.e. } t \in]0, T[, \\ h(t, r) &\geq -\eta(t), & \text{for every } r \geq 1 \text{ and a.e. } t \in]0, T[, \end{aligned}$$

and

$$\frac{1}{T} \int_0^T \limsup_{r \rightarrow 0^+} h(t, r) dt < \bar{e} < \frac{1}{T} \int_0^T \liminf_{r \rightarrow +\infty} h(t, r) dt.$$

(H3) *There is a constant $\bar{\delta} > 0$ and a differentiable function $F :]0, \bar{\delta}[\rightarrow \mathbb{R}$ such that*

$$h(t, r) \leq F'(r), \quad \text{for every } r \in]0, \bar{\delta}[\text{ and a.e. } t \in]0, T[,$$

and

$$\lim_{r \rightarrow 0^+} F(r) = +\infty.$$

Then, there exists a $k_1 \geq 1$ such that, for every integer $k \geq k_1$, equation (6) has a periodic solution $x_k(t)$ with minimal period kT , which makes exactly one revolution around the origin in the period time kT . Moreover, there is a constant $C > 0$ such that, for every $k \geq k_1$

$$\frac{1}{C} < |x_k(t)| < C, \quad \text{for every } t \in \mathbb{R},$$

and, if μ_k denotes the angular momentum associated to $x_k(t)$, then

$$\lim_{k \rightarrow \infty} \mu_k = 0.$$

Let us make a brief comment on the hypotheses of Theorem 1. Assumption (H1), introduced in [12], can be interpreted as a nonresonance condition with respect to the Dancer-Fučík spectrum. Indeed, the number $(\frac{\pi}{T})^2$ is the value of the asymptote to the first curve of that spectrum. Assumption (H2) is the well-known Landesman-Lazer condition, a nonresonance condition with respect to the first eigenvalue of the differential operator. Assumption (H3) is introduced in order to control the solutions which approach the singularity, by the use of some energy estimates.

As an immediate consequence of Theorem 1, in the particular case of system (3), we have the following.

Corollary 1 *Assume that, for some negative constants c_1 and c_2 ,*

$$c_1 \leq c(t) \leq c_2 < 0, \quad \text{for a.e. } t \in \mathbb{R}. \quad (8)$$

If $\gamma \geq 2$ and $\bar{e} < 0$, then the same conclusion of Theorem 1 holds for system (3).

Our second theorem deals with periodic solutions having a large angular momentum. It is closely related to [8, Theorem 4], where the attractive case was studied.

Theorem 2 *Let the following two assumptions hold.*

$$(H4) \quad \lim_{r \rightarrow +\infty} \frac{h(t, r)}{r} = 0,$$

uniformly for almost every $t \in]0, T[$.

(H2)₊ *There exists a function $\eta \in L^1(0, T)$, with positive values, such that*

$$h(t, r) \geq -\eta(t), \quad \text{for every } r \geq 1 \text{ and a.e. } t \in]0, T[,$$

and

$$\bar{e} < \frac{1}{T} \int_0^T \liminf_{r \rightarrow +\infty} h(t, r) dt.$$

Then, there exists a $k_1 \geq 1$ such that, for every integer $k \geq k_1$, equation (6) has a periodic solution $x_k(t)$ with minimal period kT , which makes exactly one revolution around the origin in the period time kT . Moreover,

$$\lim_{k \rightarrow \infty} (\min |x_k|) = +\infty, \quad \lim_{k \rightarrow \infty} \frac{\min |x_k|}{\max |x_k|} = 1,$$

and, if μ_k denotes the angular momentum associated to $x_k(t)$, then

$$\lim_{k \rightarrow \infty} \mu_k = +\infty.$$

Notice that, in Theorem 2, we have no assumptions on the behaviour of the nonlinearity near the singularity. Indeed, (H2)₊ is a kind of Landesman-Lazer condition, but it is only assumed for large positive values of r . Assumption (H4) is a non-resonance condition which will be used to show that the angular velocity of the large-amplitude solutions can be arbitrarily small.

We have the following direct consequence of Theorem 2 concerning system (3).

Corollary 2 *If $\gamma > 1$, $\bar{e} < 0$, and (8) holds, then the same conclusion of Theorem 2 holds for system (3).*

We notice that, in [7], assuming (H2), (H3) and (H4), it was proved that there is a family of subharmonic solutions with arbitrarily large minimal periods, which oscillate radially. We thus conclude with the following.

Corollary 3 *Assume (H2), (H3) and (H4). Then, system (6) has three distinct families of subharmonic solutions, with the following distinct behaviour: one oscillates radially, one rotates with small angular momentum, and one rotates with large angular momentum and large amplitude.*

The above results should be compared with those contained in [13, 6] (see also [4, 9]), where systems of the type

$$\ddot{x} + \nabla V(x) = \mathbf{e}(t) \tag{9}$$

were considered, with $V : \mathbb{R}^N \setminus \{0\} \rightarrow \mathbb{R}$, a continuously differentiable function satisfying

$$\lim_{x \rightarrow 0} V(x) = +\infty, \tag{10}$$

and $\mathbf{e} : \mathbb{R} \rightarrow \mathbb{R}^N$, a locally integrable T -periodic vector-valued function. This type of problem is not radially symmetric, and the situation is substantially different from the one considered above. Let $\bar{\mathbf{e}}$ be the mean value of $\mathbf{e}(t)$:

$$\bar{\mathbf{e}} = \frac{1}{T} \int_0^T \mathbf{e}(t) dt.$$

The following result has been proved in [13, 6].

Theorem 3 *Assume (10) and the following conditions:*

$$\limsup_{x \rightarrow 0} \langle \nabla V(x), x \rangle < 0,$$

$$\limsup_{|x| \rightarrow \infty} |\nabla V(x)| < |\bar{\mathbf{e}}|.$$

Then, equation (9) has a T -periodic solution x , and a sequence $(x_k)_k$ of kT -periodic solutions, whose minimal periods tend to infinity.

The above theorem applies to the equation

$$\ddot{x} + \bar{c} \frac{x}{|x|^\gamma} = \mathbf{e}(t), \quad (11)$$

where $\gamma > 2$, \bar{c} is a negative constant, and $\bar{\mathbf{e}} \neq 0$. Notice that, in [13], this last assumption has been shown to be necessary for the existence of periodic solutions of (11), when $\mathbf{e}(t)$ is a small bounded function.

It should be observed that, assuming $N = 2$ in Theorem 3, the solutions could remain confined in a sector of the plane, and thus not necessarily rotate around the origin. See also [14] and [3] for related results.

2 Proof of Theorem 1: solutions with small angular momentum

Let us first clarify our assumptions. The function h is L^1 -Carathéodory, i.e.,

- $h(\cdot, r)$ is measurable, for every $r > 0$;
- $h(t, \cdot)$ is continuous, for almost every $t \in]0, T[$;
- for every compact interval $[a, b]$ in $]0, +\infty[$, there exists $\ell_{a,b} \in L^1(0, T)$ such that

$$r \in [a, b] \quad \Rightarrow \quad |h(t, r)| \leq \ell_{a,b}(t), \quad \text{for a.e. } t \in]0, T[.$$

Concerning the function $\alpha(t)$ in assumption (H1), we can assume that $\alpha(t) \geq 0$ for almost every $t \in]0, T[$. Hence, in particular, we assume $\alpha \in L^\infty(0, T)$.

In the sequel, we will implicitly assume that all functions defined (almost everywhere) on $]0, T[$ are extended by T -periodicity to the whole real line.

Let $H_0^1(0, T)$ denote the usual Sobolev space of functions satisfying the homogeneous Dirichlet boundary condition. Given a constant $\omega \in \mathbb{R}$, let $B_{\alpha, \omega} : H_0^1(0, T) \rightarrow \mathbb{R}$ be the quadratic form defined by

$$B_{\alpha, \omega}(v) = \int_0^T [\dot{v}^2(t) - \alpha(t + \omega)v^2(t)] dt.$$

We will need the following lemma, analogous to [12, Lemma 3].

Lemma 1 *There is a constant $\bar{\varepsilon} > 0$ such that*

$$B_{\alpha, \omega}(v) \geq \bar{\varepsilon} \int_0^T \dot{v}^2(t) dt,$$

for every $\omega \in \mathbb{R}$ and $v \in H_0^1(0, T)$.

Proof By (7) and the Poincaré inequality,

$$B_{\alpha, \omega}(v) \geq \int_0^T \left[\dot{v}^2(t) - \left(\frac{\pi}{T} \right)^2 v^2(t) \right] dt \geq 0, \quad (12)$$

for every $\omega \in \mathbb{R}$ and $v \in H_0^1(0, T)$. Let us see that

$$B_{\alpha, \omega}(v) = 0 \quad \Leftrightarrow \quad v = 0.$$

Indeed, by (12), $B_{\alpha, \omega}(v) = 0$ implies $v(t) = A \sin(\frac{\pi t}{T})$, for some constant $A \in \mathbb{R}$. Then,

$$\int_0^T \left[A^2 \left(\frac{\pi}{T} \right)^2 \cos^2 \left(\frac{\pi t}{T} \right) - \alpha(t + \omega) A^2 \sin^2 \left(\frac{\pi t}{T} \right) \right] dt = 0,$$

i.e., since $\int_0^T \cos^2(\frac{\pi t}{T}) dt = \int_0^T \sin^2(\frac{\pi t}{T}) dt$,

$$A^2 \int_0^T \left[\left(\frac{\pi}{T} \right)^2 - \alpha(t + \omega) \right] \sin^2 \left(\frac{\pi t}{T} \right) dt = 0,$$

which implies $A = 0$, as $\alpha(\cdot) < (\frac{\pi}{T})^2$ on a subset of positive measure.

Assume now, by contradiction, that for every integer $n \geq 1$ there are $\omega_n \in [0, T]$ and $v_n \in H_0^1(0, T)$ such that

$$\int_0^T [\dot{v}_n^2(t) - \alpha(t + \omega_n)v_n^2(t)] dt < \frac{1}{n} \int_0^T \dot{v}_n^2(t) dt.$$

Let $z_n = v_n / \|v_n\|_{H_0^1}$, where

$$\|v_n\|_{H_0^1} = \left(\int_0^T \dot{v}_n^2(t) dt \right)^{1/2}$$

is the usual norm in $H_0^1(0, T)$. Then, $z_n \in H_0^1(0, T)$, $\|z_n\|_{H_0^1} = 1$, and

$$\int_0^T \alpha(t + \omega_n) z_n^2(t) dt > 1 - \frac{1}{n}.$$

Passing to subsequences, we can assume that $\omega_n \rightarrow \omega \in [0, T]$, $z_n \rightharpoonup z$ (weakly) in $H_0^1(0, T)$, and $z_n \rightarrow z$ uniformly. Then, $\|z\|_{H_0^1} \leq 1$, and, since by Lebesgue's Theorem $\alpha(\cdot + \omega_n) \rightarrow \alpha(\cdot + \omega)$ in $L^1(0, T)$,

$$\int_0^T \alpha(t + \omega) z^2(t) dt \geq 1. \quad (13)$$

Hence, $B_{\alpha, \omega}(z) \leq 0$, so that, by the above, it has to be $z = 0$. We thus get a contradiction with (13). \blacksquare

Let us now begin the proof of Theorem 1. We may write the solutions of (6) in polar coordinates:

$$x(t) = \rho(t)(\cos \varphi(t), \sin \varphi(t)), \quad (14)$$

and (2) is satisfied if $\rho(t) > 0$, for every t . Equation (6) is then equivalent to the system

$$(S) \quad \begin{cases} \ddot{\rho} - \frac{\mu^2}{\rho^3} + h(t, \rho) = e(t), \\ \rho^2 \dot{\varphi} = \mu, \end{cases}$$

where μ is the (scalar) angular momentum of $x(t)$. Recall that μ is constant in time along any solution. In the following, when considering a solution of (S), we will always implicitly assume that $\rho > 0$.

Without loss of generality we assume that $e(t)$ has zero mean value, i.e.,

$$\bar{e} = 0. \quad (15)$$

Indeed, otherwise, we just replace $e(t)$ by $e(t) - \bar{e}$ and $h(t, \rho)$ by $h(t, \rho) - \bar{e}$. We consider $\mu \geq 0$ as a parameter, and, by the use of degree theory, we will prove the existence of a T -periodic solution ρ of the first equation in (S). To this aim, for $\lambda \in [0, 1]$, we introduce the modified problem

$$\begin{cases} \ddot{\rho} - \frac{\mu^2}{\rho^3} + (1 - \lambda)\left(1 - \frac{1}{\rho^3}\right) + \lambda h(t, \rho) = \lambda e(t), \\ \rho(0) = \rho(T), \quad \dot{\rho}(0) = \dot{\rho}(T). \end{cases} \quad (16)$$

For some $r_0 \in]0, 1[$, to be fixed later, we define the truncation $g_{\lambda, \mu, r_0} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, as follows:

$$g_{\lambda, \mu, r_0}(t, r) = \begin{cases} -\frac{\mu^2}{r^3} + (1 - \lambda)\left(1 - \frac{1}{r^3}\right) + \lambda h(t, r) & \text{for } r \geq r_0 \\ -\frac{\mu^2}{r_0^3} + (1 - \lambda)\left(1 - \frac{1}{r_0^3}\right) + \lambda h(t, r_0) & \text{for } r \leq r_0. \end{cases}$$

This function is still L^1 -Carathéodory. We consider the T -periodic problem

$$(P_{\lambda,\mu,r_0}) \quad \begin{cases} \ddot{\rho} + g_{\lambda,\mu,r_0}(t, \rho) = \lambda e(t), \\ \rho(0) = \rho(T), \quad \dot{\rho}(0) = \dot{\rho}(T), \end{cases}$$

and look for a priori bounds for the solutions ρ , for small values of r_0 .

Lemma 2 *Assume (H1)–(H3). Given $M > 0$, there exist $\bar{r}_0 > 0$ and $C > 0$ such that, if $\rho(t)$ is a solution of (P_{λ,μ,r_0}) , with $\lambda \in [0, 1]$, $\mu \in [0, M]$ and $r_0 \in]0, \bar{r}_0]$, then*

$$\frac{1}{C} < \rho(t) < C \quad \text{and} \quad |\dot{\rho}(t)| < C,$$

for every $t \in [0, T]$.

Proof By contradiction, assume that, for every $n \geq 1$, there are $\lambda_n \in [0, 1]$, $\mu_n \in [0, M]$, $r_{0,n} \in]0, \frac{1}{n}[$, and a solution $\rho_n(t)$ of $(P_{\lambda_n,\mu_n,r_{0,n}})$, such that, either $\rho_n([0, T]) \not\subseteq]\frac{1}{n}, n[$, or $\|\dot{\rho}_n\|_{L^\infty} \geq n$. For simplicity we denote by g_n the function $g_{\lambda_n,\mu_n,r_{0,n}}$. The remaining of the proof is divided into five steps.

Step 1. There exists $R_1 \geq 1$ such that $\min \rho_n \leq R_1$, for every n .

Otherwise, there would exist a subsequence such that $\min \rho_n \rightarrow +\infty$. We may also assume that $\lambda_n \rightarrow \bar{\lambda} \in [0, 1]$. Integrating the equation

$$\ddot{\rho}_n + g_n(t, \rho_n) = \lambda_n e(t), \tag{17}$$

by (15) we have

$$\int_0^T g_n(t, \rho_n(t)) dt = 0. \tag{18}$$

Hence, from the definition of g_n , since $\min \rho_n \rightarrow +\infty$,

$$(1 - \bar{\lambda})T + \lim_n \int_0^T \lambda_n h(t, \rho_n(t)) dt = 0.$$

Using Fatou's Lemma,

$$(1 - \bar{\lambda})T + \bar{\lambda} \int_0^T \liminf_{r \rightarrow +\infty} h(t, r) dt \leq 0,$$

and we get a contradiction with (H2).

Step 2. There exists $R_2 > R_1$ such that $\max \rho_n \leq R_2$, for every n .

Let $\bar{\varepsilon} > 0$ be as in Lemma 1. By (H1), there is a $\tilde{R}_1 \geq R_1$ such that

$$h(t, r) \leq \left(\alpha(t) + \frac{\bar{\varepsilon}}{2} \left(\frac{\pi}{T} \right)^2 \right) r, \quad \text{for every } r \geq \tilde{R}_1 \text{ and a.e. } t \in \mathbb{R}. \tag{19}$$

By contradiction, we assume that $\max \rho_n \rightarrow +\infty$. Then, for n large enough, there exists an interval $[a_n, b_n]$, with $b_n - a_n \leq T$, such that

$$\rho_n(a_n) = \tilde{R}_1 = \rho_n(b_n),$$

and

$$\rho_n(t) > \tilde{R}_1, \quad \text{for every } t \in]a_n, b_n[,$$

and there exists $t_n \in]a_n, b_n[$ such that $\max \rho_n = \rho_n(t_n)$. Let $u_n(t) = \rho_n(t) - \tilde{R}_1$, so that $u_n(a_n) = 0 = u_n(b_n)$. Recall the Poincaré inequality

$$\int_{a_n}^{b_n} u_n^2(t) dt \leq \left(\frac{b_n - a_n}{\pi} \right)^2 \int_{a_n}^{b_n} \dot{u}_n^2(t) dt. \quad (20)$$

Define $v_n : [0, T] \rightarrow \mathbb{R}$ as follows:

$$v_n(t) = \begin{cases} u_n(t + a_n) & \text{if } t \in [0, b_n - a_n], \\ 0 & \text{otherwise.} \end{cases}$$

Then, $v_n \in H_0^1(0, T)$ and, by Lemma 1,

$$B_{\alpha, a_n}(v_n) \geq \bar{\varepsilon} \int_0^T \dot{v}_n^2(t) dt,$$

for every n , i.e.,

$$\int_{a_n}^{b_n} [\dot{u}_n^2(t) - \alpha(t) u_n^2(t)] dt \geq \bar{\varepsilon} \int_{a_n}^{b_n} \dot{u}_n^2(t) dt. \quad (21)$$

Multiplying by u_n in equation (17) and integrating between a_n and b_n , by (19) we obtain

$$\begin{aligned} \int_{a_n}^{b_n} \dot{u}_n^2 &= \int_{a_n}^{b_n} u_n [g_n(t, \rho_n) - \lambda_n e(t)] dt \\ &\leq \int_{a_n}^{b_n} u_n [1 + \lambda_n (h(t, \rho_n) - e(t))] dt \\ &\leq \int_{a_n}^{b_n} u_n \left[1 + \left(\alpha(t) + \frac{\bar{\varepsilon}}{2} \left(\frac{\pi}{T} \right)^2 \right) (u_n + \tilde{R}_1) + |e(t)| \right] dt \\ &\leq \int_{a_n}^{b_n} \left[\left(\alpha(t) + \frac{\bar{\varepsilon}}{2} \left(\frac{\pi}{T} \right)^2 \right) u_n^2 + \gamma(t) u_n \right] dt, \end{aligned}$$

where

$$\gamma(t) = \tilde{R}_1 \left(\alpha(t) + \frac{\bar{\varepsilon}}{2} \left(\frac{\pi}{T} \right)^2 \right) + |e(t)| + 1.$$

Then, using (21),

$$\bar{\varepsilon} \int_{a_n}^{b_n} \dot{u}_n^2 \leq \frac{\bar{\varepsilon}}{2} \left(\frac{\pi}{T} \right)^2 \int_{a_n}^{b_n} u_n^2 + \|\gamma\|_{L^2} \left(\int_{a_n}^{b_n} u_n^2 \right)^{\frac{1}{2}}$$

so that, by (20), as $b_n - a_n \leq T$, the sequence $\left(\int_{a_n}^{b_n} \dot{u}_n^2\right)_n$ has to be bounded. Since

$$\max u_n \leq \int_{a_n}^{b_n} |\dot{u}_n| \leq \sqrt{T} \left(\int_{a_n}^{b_n} \dot{u}_n^2\right)^{\frac{1}{2}},$$

we get a contradiction with the fact that $\max \rho_n \rightarrow +\infty$.

Step 3. There exists $r_1 \in]0, 1[$ such that $\max \rho_n \geq r_1$, for every n .

Otherwise, there would exist a subsequence such that $\max \rho_n \rightarrow \bar{\rho} \in [-\infty, 0]$. We may also assume that $\lambda_n \rightarrow \bar{\lambda} \in [0, 1]$ and $\mu_n \rightarrow \bar{\mu} \in [0, M]$. Set

$$\tilde{\rho}_n(t) = \max\{\rho_n(t), r_{0,n}\}. \quad (22)$$

Notice that

$$g_n(t, \rho_n(t)) = g_n(t, \tilde{\rho}_n(t)),$$

for almost every t . Since $r_{0,n} \leq \tilde{\rho}_n(t) \leq 1$ for n large enough, by (H2),

$$\begin{aligned} g_n(t, \rho_n(t)) &= -\frac{\mu_n^2}{\tilde{\rho}_n^3(t)} + (1 - \lambda_n) \left(1 - \frac{1}{\tilde{\rho}_n^3(t)}\right) + \lambda_n h(t, \tilde{\rho}_n(t)) \\ &\leq 1 + \lambda_n h(t, \tilde{\rho}_n(t)) \\ &\leq 1 + \eta(t). \end{aligned}$$

Hence, by Fatou's Lemma and (18), we have

$$\int_0^T \limsup_n g_n(t, \rho_n(t)) dt \geq \limsup_n \int_0^T g_n(t, \rho_n(t)) dt = 0,$$

and then

$$\int_0^T \limsup_n \left[(1 - \lambda_n) \left(1 - \frac{1}{\tilde{\rho}_n^3(t)}\right) + \lambda_n h(t, \tilde{\rho}_n(t)) \right] dt \geq 0.$$

Since $\tilde{\rho}_n(t) > 0$ for every t and $\max \tilde{\rho}_n(t) \rightarrow 0$, we easily get a contradiction with (H2).

Step 4. Proof of the estimate on the derivative.

By (H2), we have

$$g_n(t, r) \leq \eta(t), \quad \text{for every } r \leq 1 \text{ and a.e. } t \in]0, T[.$$

Then, using (18),

$$\begin{aligned} \int_{\{\rho_n \leq 1\}} |g_n(t, \rho_n(t))| dt &\leq \int_{\{\rho_n \leq 1\}} |g_n(t, \rho_n(t)) - \eta(t)| dt + \|\eta\|_{L^1} \\ &= \int_{\{\rho_n \leq 1\}} (-g_n(t, \rho_n(t)) + \eta(t)) dt + \|\eta\|_{L^1} \\ &\leq - \int_{\{\rho_n \leq 1\}} g_n(t, \rho_n(t)) dt + 2\|\eta\|_{L^1} \\ &= \int_{\{\rho_n > 1\}} g_n(t, \rho_n(t)) dt + 2\|\eta\|_{L^1} \\ &\leq \int_{\{\rho_n > 1\}} |g_n(t, \rho_n(t))| dt + 2\|\eta\|_{L^1}, \end{aligned}$$

so that

$$\begin{aligned} \int_0^T |g_n(t, \rho_n(t))| dt &= \left(\int_{\{\rho_n \leq 1\}} + \int_{\{\rho_n > 1\}} \right) |g_n(t, \rho_n(t))| dt \\ &\leq 2 \int_{\{\rho_n > 1\}} |g_n(t, \rho_n(t))| dt + 2\|\eta\|_{L^1}. \end{aligned}$$

As proved in Step 2, we have $\rho_n(t) \leq R_2$, for every t . So, there is a constant $c_1 > 0$ for which

$$\int_{\{\rho_n > 1\}} |g_n(t, \rho_n(t))| dt \leq c_1,$$

for every n . Consequently,

$$\int_0^T |\ddot{\rho}_n(t)| dt \leq \int_0^T |g_n(t, \rho_n(t))| dt + \|e\|_{L^1} \leq 2c_1 + 2\|\eta\|_{L^1} + \|e\|_{L^1}.$$

Since, being ρ_n periodic, its derivative must vanish somewhere, we have that $\|\dot{\rho}_n\|_{L^\infty} \leq \|\ddot{\rho}_n\|_{L^1}$. Setting $C_1 = 2c_1 + 2\|\eta\|_{L^1} + \|e\|_{L^1}$, we thus have

$$\|\dot{\rho}_n\|_{L^\infty} \leq C_1, \quad (23)$$

for every n .

Step 5. Conclusion of the proof.

Using Step 2 and Step 4, if n is sufficiently large we have that $\rho_n(t) < n$ and $\|\dot{\rho}_n(t)\| < n$, for every t . Therefore, it has to be $\min \rho_n \leq \frac{1}{n}$, for n large enough. Let $r_1 \in]0, 1[$ be as in Step 3, and set $\tilde{r}_1 = \min\{r_1, \bar{\delta}\}$. We can assume $\tilde{r}_1 > \frac{1}{n}$. Then, there is an interval $[\gamma_n, \delta_n]$ such that

$$\rho_n(\gamma_n) = \tilde{r}_1, \quad \rho_n(\delta_n) = \frac{1}{n},$$

and

$$\frac{1}{n} < \rho_n(t) < \tilde{r}_1, \quad \text{for every } t \in]\gamma_n, \delta_n[.$$

Recall that $r_{0,n} < \frac{1}{n}$. So, for $t \in [\gamma_n, \delta_n]$, we have that $\rho_n(t) = \tilde{\rho}_n(t)$. We define

$$\tilde{h}(t, r) = h(t, r) - \eta(t), \quad \tilde{e}(t) = e(t) - \eta(t), \quad \tilde{f}(r) = \min\{F'(r), 0\},$$

and let $\tilde{F} :]0, \bar{\delta}[\rightarrow \mathbb{R}$ be a primitive of \tilde{f} , i.e., $\tilde{F}'(r) = \tilde{f}(r)$, for every r . Then,

$$\tilde{h}(t, r) \leq \tilde{F}'(r), \quad \text{for every } r \in]0, \bar{\delta}[\text{ and a.e. } t \in \mathbb{R},$$

and

$$\lim_{r \rightarrow 0^+} \tilde{F}(r) = +\infty. \quad (24)$$

We can write (17) as

$$\ddot{\rho}_n + \tilde{g}_n(t, \rho_n) = \lambda_n \tilde{e}(t), \quad (25)$$

where $\tilde{g}_n(t, r)$ is defined as $g_n(t, r)$, with $\tilde{h}(t, r)$ instead of $h(t, r)$. Notice that

$$\tilde{g}_n(t, \rho_n(t)) = -\frac{\mu_n^2}{\rho_n^3(t)} + (1 - \lambda_n) \left(1 - \frac{1}{\rho_n^3(t)}\right) + \lambda_n \tilde{h}(t, \rho_n(t)) \leq 0,$$

for every $t \in]\gamma_n, \delta_n[$. Let C_1 be the constant defined in Step 4, for which (23) holds. Multiplying in (25) by $(\dot{\rho}_n - C_1)$ and integrating on $[\gamma_n, \delta_n]$,

$$\begin{aligned} \frac{1}{2}[(\dot{\rho}_n - C_1)^2]_{\gamma_n}^{\delta_n} + \int_{\gamma_n}^{\delta_n} \tilde{g}_n(t, \rho_n)(\dot{\rho}_n - C_1) dt &= \int_{\gamma_n}^{\delta_n} \lambda_n \tilde{e}(t)(\dot{\rho}_n - C_1) dt \\ &\leq 2C_1(\|e\|_{L^1} + \|\eta\|_{L^1}). \end{aligned}$$

Hence, there is a constant $c_2 > 0$ for which

$$\int_{\gamma_n}^{\delta_n} \tilde{g}_n(t, \rho_n)(\dot{\rho}_n - C_1) dt \leq c_2, \quad (26)$$

for every n . On the other hand, since $\dot{\rho}_n - C_1 \leq 0$,

$$\begin{aligned} \int_{\gamma_n}^{\delta_n} \tilde{g}_n(t, \rho_n)(\dot{\rho}_n - C_1) dt &\geq \\ &\geq \int_{\gamma_n}^{\delta_n} \left((1 - \lambda_n) \left(1 - \frac{1}{\rho_n^3}\right) + \lambda_n \tilde{h}(t, \rho_n) \right) (\dot{\rho}_n - C_1) dt \\ &\geq \int_{\gamma_n}^{\delta_n} \left((1 - \lambda_n) \left(1 - \frac{1}{\rho_n^3}\right) + \lambda_n \tilde{F}'(\rho_n) \right) (\dot{\rho}_n - C_1) dt \\ &\geq \int_{\gamma_n}^{\delta_n} \left((1 - \lambda_n) \left(1 - \frac{1}{\rho_n^3}\right) + \lambda_n \tilde{F}'(\rho_n) \right) \dot{\rho}_n dt \\ &= \left[(1 - \lambda_n) \left(\rho_n + \frac{2}{\rho_n^2} \right) + \lambda_n \tilde{F}(\rho_n) \right]_{\gamma_n}^{\delta_n} \\ &\geq (1 - \lambda_n) \left(\frac{1}{n} + 2n^2 \right) + \lambda_n \tilde{F}\left(\frac{1}{n}\right) - c_3, \end{aligned}$$

for some constant $c_3 > 0$. Using (24), when n tends to infinity we get a contradiction with (26), thus ending the proof of the lemma. \blacksquare

We now fix $M > 0$ and take $\bar{r}_0 > 0$ and $C > 0$ as given by Lemma 2. Let us also fix $r_0 = \min\{\bar{r}_0, \frac{1}{C}\}$. As a consequence of Lemma 2, if $\lambda \in [0, 1]$ and $\mu \in [0, M]$, any solution $\rho(t)$ of (P_{λ, μ, r_0}) is also a solution of (16). In particular, if $\lambda = 1$, any solution of (P_{1, μ, r_0}) is a T -periodic solution of the first equation in (S).

Notice that, viceversa, once $M > 0$ is fixed, every T -periodic solution of the first equation in (S) with $\mu \in [0, M]$ satisfies (P_{1, μ, \hat{r}_0}) for sufficiently small $\hat{r}_0 > 0$, so that it also verifies the estimates given by Lemma 2.

In the following, we denote by C_T^1 the set of T -periodic C^1 -functions, with the usual norm of $C^1([0, T])$.

Lemma 3 *Given $M > 0$, there is a continuum \mathcal{C} in $[0, M] \times C_T^1$, connecting $\{0\} \times C_T^1$ with $\{M\} \times C_T^1$, whose elements (μ, ρ) satisfy the first equation in (S).*

Proof In order to apply degree theory, let us define the following operators:

$$\begin{aligned} L : D(L) &\subset C^1([0, T]) \rightarrow L^1(0, T), \\ D(L) &= \{\rho \in W^{2,1}(0, T) : \rho(0) = \rho(T), \dot{\rho}(0) = \dot{\rho}(T)\}, \\ L\rho &= \ddot{\rho} - \rho, \end{aligned}$$

and, for $\lambda \in [0, 1]$,

$$\begin{aligned} N_\lambda : [0, M] \times C^1([0, T]) &\rightarrow L^1(0, T), \\ N_\lambda(\mu, \rho)(t) &= -g_{\lambda, \mu, r_0}(t, \rho(t)) + \lambda e(t) - \rho(t). \end{aligned}$$

Problem (P_{λ, μ, r_0}) is thus equivalent to

$$L\rho = N_\lambda(\mu, \rho).$$

Since L is invertible, we can write equivalently

$$\rho - L^{-1}N_\lambda(\mu, \rho) = 0. \quad (27)$$

Let $C > 0$ be the constant given by Lemma 2 and define Ω to be the following open subset of $C^1([0, T])$:

$$\Omega = \left\{ \rho \in C^1([0, T]) : \frac{1}{C} < \rho(t) < C \text{ and } |\dot{\rho}(t)| < C, \text{ for every } t \in [0, T] \right\}.$$

By Lemma 2, equation (27) has no solutions ρ on $\partial\Omega$, for any $\lambda \in [0, 1]$ and $\mu \in [0, M]$. Since $L^{-1}N_\lambda(\mu, \cdot)$ is a compact operator, by the global continuation principle of Leray-Schauder (see e.g. [15, Theorem 14.C]), the lemma will be proved if we show that the degree is nonzero for some $(\lambda, \mu) \in [0, 1] \times [0, M]$. Indeed, since the degree is the same for every $(\lambda, \mu) \in [0, 1] \times [0, M]$, then, it will be nonzero when $\lambda = 1$, for every $\mu \in [0, M]$.

Let us then take $\mu = \lambda = 0$ and define the function

$$\begin{aligned} \Psi : \left[\frac{1}{C}, C \right] \times [-C, C] &\rightarrow \mathbb{R}^2, \\ \Psi(u, v) &= \left(v, \frac{1}{u^3} - 1 \right). \end{aligned}$$

By a result of Capietto, Mawhin and Zanolin [2, Theorem 1], one can compute the Leray-Schauder degree of $I - L^{-1}N_0(0, \cdot)$ as the Brouwer degree of Ψ :

$$d_{LS}(I - L^{-1}N_0(0, \cdot), \Omega) = d_B\left(\Psi, \left] \frac{1}{C}, C \right[\times] - C, C \right).$$

Since Ψ has the unique zero $(1, 0)$, and the jacobian matrix $J_\Psi(1, 0)$ has a positive determinant, we conclude that the degree has to be equal to 1. \blacksquare

Notice that, if $(\mu, \rho) \in \mathcal{C}$ then ρ is T -periodic and the first equation in (S) is satisfied. Let us consider the function $\Phi : \mathcal{C} \rightarrow \mathbb{R}$, defined by

$$\Phi(\mu, \rho) \mapsto \int_0^T \frac{\mu}{\rho^2(t)} dt.$$

It is continuous and defined on a connected domain, so its image is an interval. Since $\Phi(0, \rho) = 0$ and, by Lemma 2,

$$\Phi(\mu, \rho) \leq TC^2\mu,$$

this interval is of the type $[0, \bar{\theta}]$ for some $\bar{\theta} > 0$.

Lemma 4 *For every $\theta \in [0, \bar{\theta}]$, there are (μ, ρ, φ) , verifying system (S), for which $(\mu, \rho) \in \mathcal{C}$, and*

$$\rho(t+T) = \rho(t), \quad \varphi(t+T) = \varphi(t) + \theta,$$

for every $t \in \mathbb{R}$.

Proof Given $\theta \in [0, \bar{\theta}]$, there are $(\mu, \rho) \in \mathcal{C}$ such that

$$\int_0^T \frac{\mu}{\rho^2(t)} dt = \theta.$$

As noticed above, the first equation in (S) is satisfied. Moreover, defining

$$\varphi(t) = \int_0^t \frac{\mu}{\rho^2(s)} ds,$$

the second equation in (S) is also satisfied and

$$\varphi(t+T) - \varphi(t) = \int_t^{t+T} \frac{\mu}{\rho^2(s)} ds = \int_0^T \frac{\mu}{\rho^2(s)} ds = \theta.$$

■

We are going to complete now the proof of Theorem 1. For every $\theta \in [0, \bar{\theta}]$, the solution of system (S) found in Lemma 4 provides, through (14), a solution to equation (6) such that

$$x(t+T) = e^{i\theta}x(t),$$

for every $t \in \mathbb{R}$ (for briefness we used here the complex notation).

In particular, if $\theta = \frac{2\pi}{k}$ for some integer $k \geq 1$, then $x(t)$ is periodic with minimal period kT , and rotates exactly once around the origin in the period time kT . Hence, for every integer $k \geq 2\pi/\bar{\theta}$, we have such a kT -periodic solution, which we denote by $x_k(t)$. Let $(\rho_k(t), \varphi_k(t))$ be its polar coordinates,

and μ_k be its angular momentum. By the above construction, $(\mu_k, \rho_k, \varphi_k)$ verify system (S), $(\mu_k, \rho_k) \in \mathcal{C}$, and

$$\int_0^T \frac{\mu_k}{\rho_k^2(t)} dt = \frac{2\pi}{k}.$$

Since $\mu_k \in [0, M]$, by Lemma 2 we have that

$$\frac{1}{C} < \rho_k(t) < C \quad \text{and} \quad |\dot{\rho}_k(t)| < C,$$

for every $t \in [0, T]$. Hence,

$$\frac{2\pi}{k} = \int_0^T \frac{\mu_k}{\rho_k^2(t)} dt > T \frac{\mu_k}{C^2},$$

so that

$$\lim_{k \rightarrow \infty} \mu_k = 0.$$

The proof is thus completed. ■

3 Proof of Theorem 2: solutions with large angular momentum

The proof of Theorem 2 follows closely both the proof of [8, Theorem 4], where the attractive case has been considered, and the proof of Theorem 1.

As in the previous section, without loss of generality, we assume (15), i.e.,

$$\bar{e} = 0.$$

We consider $\mu \geq 0$ as a parameter, and, by the use of degree theory, we will prove the existence of a T -periodic solution ρ of the first equation in (S). To this aim, for $\lambda \in [0, 1]$, we introduce the modified problem (16). We define the truncation at $r_0 = 1$, i.e., $g_{\lambda, \mu} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, as follows:

$$g_{\lambda, \mu}(t, r) = \begin{cases} -\frac{\mu^2}{r^3} + (1 - \lambda)\left(1 - \frac{1}{r^3}\right) + \lambda h(t, r) & \text{for } r \geq 1 \\ -\mu^2 + \lambda h(t, 1) & \text{for } r \leq 1. \end{cases}$$

This function is still L^1 -Carathéodory. We consider the T -periodic problem

$$(P_{\lambda, \mu}) \quad \begin{cases} \ddot{\rho} + g_{\lambda, \mu}(t, \rho) = \lambda e(t), \\ \rho(0) = \rho(T), \quad \dot{\rho}(0) = \dot{\rho}(T), \end{cases}$$

and look for a priori bounds for the positive solutions ρ . This time, we will consider solutions with a large angular momentum.

Lemma 5 *For every $\Gamma > 1$ there exists $\mu(\Gamma) \geq 1$ such that, if $\mu \geq \mu(\Gamma)$, $\lambda \in [0, 1]$, and ρ is a positive solution of $(P_{\lambda, \mu})$, then $\|\rho\|_{L^\infty} \geq \Gamma$.*

Proof Assume by contradiction that there are $\Gamma > 1$, and sequences $(\lambda_n)_n$, $(\mu_n)_n$, and $(\rho_n)_n$ such that $\lambda_n \in [0, 1]$, $\lim_n \mu_n = +\infty$, and ρ_n is a positive solution of (P_{λ_n, μ_n}) , with $\|\rho_n\|_{L^\infty} < \Gamma$. Integrating the equation, since $\bar{e} = 0$, we obtain

$$\int_0^T g_{\lambda_n, \mu_n}(t, \rho_n(t)) dt = 0.$$

Hence, setting $\tilde{\rho}_n(t) = \max\{\rho_n(t), 1\}$,

$$\mu_n^2 \int_0^T \frac{1}{\tilde{\rho}_n^3(t)} dt = (1 - \lambda_n) \int_0^T \left(1 - \frac{1}{\tilde{\rho}_n^3(t)}\right) dt + \lambda_n \int_0^T h(t, \tilde{\rho}_n(t)) dt.$$

Since $1 \leq \tilde{\rho}_n \leq \Gamma$, using the L^1 -Carathéodory condition we see that the right-hand side is bounded, and we get a contradiction with the assumption that $(\mu_n)_n$ tends to $+\infty$. \blacksquare

Lemma 6 *There exists a constant $\bar{C} > 0$ such that, if $\mu \geq 1$, $\lambda \in [0, 1]$, and ρ is a positive solution of $(P_{\lambda, \mu})$, then*

$$\min \rho \geq \frac{1}{2} \|\rho\|_{L^\infty} - \bar{C}.$$

Proof By (H4), we can fix $\bar{r} \geq 1$ such that

$$r \geq \bar{r} \quad \Rightarrow \quad |h(t, r)| \leq \frac{r}{4T^2} \quad \text{for a.e. } t \in]0, T[.$$

Multiplying in $(P_{\lambda, \mu})$ by ρ and integrating we get, writing $\tilde{\rho} = \max\{\rho, 1\}$,

$$\begin{aligned} \int_0^T \dot{\rho}^2 &= - \int_0^T \frac{\mu^2}{\tilde{\rho}^3} \rho + (1 - \lambda) \int_0^T \left(1 - \frac{1}{\tilde{\rho}^3}\right) \rho + \lambda \int_0^T h(t, \tilde{\rho}) \rho - \lambda \int_0^T e \rho \\ &\leq T \|\rho\|_{L^\infty} + \int_{\{1 \leq \tilde{\rho} < \bar{r}\}} |h(t, \tilde{\rho})| \rho + \int_{\{\tilde{\rho} \geq \bar{r}\}} |h(t, \tilde{\rho})| \rho + \|e\|_{L^1} \|\rho\|_{L^\infty} \\ &\leq \frac{1}{4T} \|\rho\|_{L^\infty}^2 + (T + \|\ell_{1, \bar{r}}\|_{L^1} + \|e\|_{L^1}) \|\rho\|_{L^\infty}. \end{aligned}$$

Setting

$$\bar{C} := (T + \|\ell_{1, \bar{r}}\|_{L^1} + \|e\|_{L^1})T,$$

we have

$$T \|\dot{\rho}\|_{L^2}^2 \leq \frac{1}{4} (\|\rho\|_{L^\infty}^2 + 4\bar{C} \|\rho\|_{L^\infty}) \leq \frac{1}{4} (\|\rho\|_{L^\infty} + 2\bar{C})^2.$$

So,

$$\max \rho - \min \rho \leq \sqrt{T} \|\dot{\rho}\|_{L^2} \leq \frac{1}{2} \|\rho\|_{L^\infty} + \bar{C},$$

thus proving the lemma. \blacksquare

Let us now fix $\bar{\Gamma} > 2(1 + \bar{C})$, where \bar{C} is given by Lemma 6. Correspondingly, let $\bar{\mu} := \mu(\bar{\Gamma})$, with $\mu(\bar{\Gamma}) \geq 1$ as in Lemma 5.

Lemma 7 *Given A, B , with $\bar{\mu} \leq A \leq B$, there is a constant $C > 1$ such that, if $\mu \in [A, B]$, $\lambda \in [0, 1]$, and ρ is a positive solution of $(P_{\lambda, \mu})$, then*

$$1 < \rho(t) < C, \quad |\dot{\rho}(t)| < C,$$

for every $t \in [0, T]$.

Proof By contradiction, assume that, for every $n \geq 1$ there are $\lambda_n \in [0, 1]$, $\mu_n \in [A, B]$ and a positive solution $\rho_n(t)$ of the T -periodic problem (P_{λ_n, μ_n}) such that, either $\rho_n([0, T]) \not\subseteq]1, n[$ or $\|\dot{\rho}_n\|_{L^\infty} \geq n$. By the choice of $\bar{\mu}$ and Lemma 5, it has to be $\|\rho_n\|_{L^\infty} \geq \bar{\Gamma}$, and by Lemma 6,

$$\min \rho_n \geq \frac{1}{2} \|\rho_n\|_{L^\infty} - \bar{C} \geq \frac{1}{2} \bar{\Gamma} - \bar{C} > 1.$$

Arguing as in Step 1 of the proof of Lemma 2 we show that there exists a $R_1 \geq 1$ such that $\min \rho_n \leq R_1$ for every n . By Lemma 6 we deduce that

$$\|\rho_n\|_{L^\infty} \leq 2(\min \rho_n + \bar{C}) < 2(R_1 + \bar{C}).$$

Set $b = 2(R_1 + \bar{C})$. From the equation in (P_{λ_n, μ_n}) and the Carathéodory conditions, recalling that $\rho_n(t) > 1$, since $\dot{\rho}_n(t)$ vanishes somewhere, we obtain

$$\|\dot{\rho}_n\|_{L^\infty} \leq \|\ddot{\rho}_n\|_{L^1} \leq \int_0^T \frac{\mu_n^2}{\rho_n^3} + \int_0^T |h(t, \rho_n)| + \|e\|_{L^1} < B^2 T + \|\ell_{1,b}\|_{L^1} + \|e\|_{L^1},$$

thus arriving at a contradiction. \blacksquare

As a consequence of Lemma 7, if $\mu \geq \bar{\mu}$ and $\rho(t)$ is a T -periodic solution of $(P_{1, \mu})$ then $\rho(t)$ also satisfies the first equation in (S). The following lemma gives an important information concerning the T -periodic solutions of that equation.

Lemma 8 *Given A, B , with $\bar{\mu} \leq A \leq B$, there is a continuum $\mathcal{C}_{A,B}$ in $[A, B] \times C_T^1$, connecting $\{A\} \times C_T^1$ with $\{B\} \times C_T^1$, whose elements (μ, ρ) satisfy the first equation in (S), being $\rho(t) > 1$ for every t .*

Proof We proceed as in the proof of Lemma 3. Let L be the differential operator defined there, and, for $\lambda \in [0, 1]$,

$$\begin{aligned} N_\lambda &: [A, B] \times C^1([0, T]) \rightarrow L^1(0, T), \\ N_\lambda(\mu, \rho)(t) &= -g_{\lambda, \mu}(t, \rho(t)) + \lambda e(t) - \rho(t). \end{aligned}$$

Problem $(P_{\lambda, \mu})$ is thus equivalent to (27). Let $C > 0$ be the constant given by Lemma 7 and define Ω to be the following open subset of $C^1([0, T])$:

$$\Omega = \left\{ \rho \in C^1([0, T]) : 1 < \rho(t) < C \text{ and } |\dot{\rho}(t)| < C, \text{ for every } t \in [0, T] \right\}.$$

By Lemma 7, equation (27) has no solutions ρ on $\partial\Omega$, for any $\lambda \in [0, 1]$ and $\mu \in [A, B]$. Since $L^{-1}N_\lambda(\mu, \cdot)$ is a compact operator, by the global continuation principle of Leray-Schauder, the lemma will be proved if we show that the degree is nonzero for some $(\lambda, \mu) \in [0, 1] \times [A, B]$.

Let us then take $\lambda = 0$, $\mu = A$, and define the function

$$\begin{aligned}\Psi &: [1, C] \times [-C, C] \rightarrow \mathbb{R}^2, \\ \Psi(u, v) &= \left(v, \frac{A^2 + 1}{u^3} - 1\right).\end{aligned}$$

Since

$$d_B\left(\Psi,]1, C[\times]-C, C[\right) = 1,$$

we conclude as in the proof of Lemma 3. ■

Using classical arguments from the theory of global continuation, we can deduce from Lemmas 7 and 8 that there is a connected set \mathcal{C} , contained in $[\bar{\mu}, +\infty[\times C_T^1$, which connects $\{\bar{\mu}\} \times C_T^1$ with $\{\mu^*\} \times C_T^1$, for every $\mu^* > \bar{\mu}$, whose elements (μ, ρ) satisfy the first equation in (S).

Lemma 9 *For every $\varepsilon > 0$, there exists $\mu_\varepsilon \geq \bar{\mu}$ such that, if $(\mu, \rho) \in \mathcal{C}$ with $\mu \geq \mu_\varepsilon$, then*

$$\int_0^T \frac{\mu}{\rho^2(t)} dt \leq \varepsilon.$$

Proof Given $\varepsilon > 0$, set

$$\varepsilon' := \left(\frac{\varepsilon}{36T}\right)^2.$$

Let \bar{C} be as in Lemma 6. By (H4), there exists $r' \geq \max\{3\bar{C}, 1\}$ such that

$$r \geq r' \quad \Rightarrow \quad |h(t, r)| \leq \varepsilon' r, \quad \text{for a.e. } t \in]0, T[.$$

For $\Gamma := 2(r' + \bar{C})$, let $\mu(\Gamma)$ be as in Lemma 5. Set $\mu_\varepsilon := \max\{\mu(\Gamma), \bar{\mu}\}$.

Let (μ, ρ) be an element of \mathcal{C} , with $\mu \geq \mu_\varepsilon$. By Lemma 5, $\|\rho\|_{L^\infty} \geq \Gamma$, and by Lemma 6, $\rho(t) \geq \frac{1}{2}\|\rho\|_{L^\infty} - \bar{C} \geq r'$, for every $t \in \mathbb{R}$. Hence,

$$\frac{1}{T} \int_0^T h(t, \rho) \leq \frac{1}{T} \int_0^T \varepsilon' \rho \leq \varepsilon' \|\rho\|_{L^\infty}.$$

Integrating the first equation in (S), since $\bar{e} = 0$, we have

$$\frac{1}{T} \int_0^T \frac{\mu^2}{\rho^3} = \frac{1}{T} \int_0^T h(t, \rho) \leq \varepsilon' \|\rho\|_{L^\infty}.$$

On the other hand,

$$\frac{1}{T} \int_0^T \frac{\mu^2}{\rho^3} \geq \frac{\mu^2}{\|\rho\|_{L^\infty}^3},$$

so that

$$\frac{\mu^2}{\|\rho\|_{L^\infty}^4} \leq \varepsilon'.$$

Then, using again Lemma 6,

$$\begin{aligned}
\int_0^T \frac{\mu}{\rho^2(t)} dt &\leq T \frac{\mu}{(\min \rho)^2} \leq T \frac{4\mu}{(\|\rho\|_{L^\infty} - 2\bar{C})^2} \\
&= T \frac{\mu}{\|\rho\|_{L^\infty}^2} \left(\frac{2\|\rho\|_{L^\infty}}{\|\rho\|_{L^\infty} - 2\bar{C}} \right)^2 \leq T \sqrt{\varepsilon'} \left(\frac{2r'}{r' - 2\bar{C}} \right)^2 \\
&\leq T \sqrt{\varepsilon'} \left(\frac{6\bar{C}}{3\bar{C} - 2\bar{C}} \right)^2 = 36 T \sqrt{\varepsilon'} = \varepsilon,
\end{aligned}$$

thus proving the lemma. ■

Since the function

$$(\mu, \rho) \mapsto \int_0^T \frac{\mu}{\rho^2(t)} dt$$

is continuous from \mathcal{C} to \mathbb{R} , and \mathcal{C} is connected, its image is an interval. By Lemmas 8 and 9 this interval is of the type $]0, \bar{\theta}]$ for some $\bar{\theta} > 0$. The analogue of Lemma 4 then holds in this case, too, with $\theta \in]0, \bar{\theta}]$.

The proof of Theorem 2 can now be completed as in the previous section (see also [8, Theorem 4]).

4 Remarks on the continuum of solutions and multiplicity of periodic solutions

Assume (H1), (H2) and (H3). By Lemma 2 we have that, for every $M > 0$, there exists a constant $C > 0$ such that, for any $\mu \in [0, M]$, if $\rho(t)$ is a T -periodic solution of the first equation in (S), then

$$\frac{1}{C} < \rho(t) < C \quad \text{and} \quad |\dot{\rho}(t)| < C,$$

for every $t \in [0, T]$. Moreover, as seen in Lemma 3, for every μ the associated degree is constantly equal to 1. Hence, by the global continuation principle of Leray-Schauder, there is a closed connected set \mathcal{C} , contained in $[0, +\infty[\times C_T^1$, which connects $\{0\} \times C_T^1$ with $\{\mu^*\} \times C_T^1$, for every $\mu^* > 0$, whose elements (μ, ρ) satisfy the first equation in (S).

Consider now the function $\hat{\varphi} : \mathcal{C} \rightarrow C^2(\mathbb{R})$, defined by

$$\hat{\varphi}(\mu, \rho)(t) = \int_0^t \frac{\mu}{\rho^2(\tau)} d\tau.$$

Since this function is continuous, its graph is a closed connected subset $\hat{\mathcal{C}}_+$ of $[0, +\infty[\times C_T^1 \times C^2(\mathbb{R})$. Moreover, the projection of $\hat{\mathcal{C}}_+$ on its first component is the whole half-line $[0, +\infty[$.

It is easily seen that the same arguments used till now symmetrically hold in the case of negative angular momenta μ . In particular, the above a priori bounds hold for every $\mu \in [-M, M]$, once $M > 0$ has been fixed. We then have the following result.

Lemma 10 *Let assumptions (H1)-(H3) hold. Then, there is a closed connected subset $\hat{\mathcal{C}}$ of $\mathbb{R} \times C_T^1 \times C^2(\mathbb{R})$ whose elements (μ, ρ, φ) solve system (S), and whose projection on its first component is the whole real line $]-\infty, +\infty[$. Moreover, for any sufficiently large integer k , this set $\hat{\mathcal{C}}$ contains solutions for which μ is small and $\varphi(t + kT) = \varphi(t) + 2\pi$, for every t .*

If, instead of (H1), the stronger assumption (H4) holds, then we can repeat the estimates made in Lemmas 5 - 9 to deduce that, along the connected set $\hat{\mathcal{C}}$ given by Lemma 10, there also are solutions with large angular momenta and amplitudes, for which $\varphi(t + kT) = \varphi(t) + 2\pi$, for every t .

Recalling that (S) represents, through

$$x(t) = \rho(t)(\cos \varphi(t), \sin \varphi(t)),$$

the solutions of (6) having angular momentum μ , we can conclude as follows.

Corollary 4 *Let assumptions (H2), (H3) and (H4) hold. Then, there is a closed connected subset of $\mathbb{R} \times C^1(\mathbb{R}, \mathbb{R}^2 \setminus \{0\})$ whose elements (μ, x) are such that $x(t)$ is a solution of system (6) with angular momentum μ , and $|x(t)|$ is T -periodic. The projection of this set on its first component is the whole real line $]-\infty, +\infty[$. Moreover, this set contains infinitely many periodic solutions, rotating around the origin, with arbitrarily large periods and amplitudes.*

Till now, we always considered periodic solutions making exactly one revolution around the origin in their period time. Concerning the existence of periodic solutions making a higher number of revolutions around the origin, we have the following result.

Theorem 4 *Let $e(t)$ be continuous, with minimal period T , and negative mean value. Let $h(t, r)$ be continuous, and assume that*

$$(H5) \quad \lim_{r \rightarrow +\infty} h(t, r) = 0,$$

uniformly for every t . Then, for every integer $m \geq 1$, there exists a $k_m \geq 1$ such that, for every integer $k \geq k_m$, if k and m are relatively prime, equation (6) has a periodic solution $x_{k,m}(t)$ with minimal period kT , which makes exactly m revolutions around the origin in the period time kT . Moreover,

$$\lim_{k \rightarrow \infty} (\min |x_{k,m}|) = +\infty,$$

and

$$\lim_{k \rightarrow \infty} \frac{\min |x_{k,m}|}{\max |x_{k,m}|} = 1.$$

We omit the proof, since it is exactly the same as that of [8, Theorem 7]. To conclude, let us state the following multiplicity result.

Theorem 5 *Let $e(t)$ be continuous, with minimal period T , and negative mean value. Let $h(t, r)$ be continuous, and assume (H5). Then, for every $n \in \mathbb{N}$, there is a $k(n) \geq 1$ such that, for every prime integer $k \geq k(n)$, equation (6) has at least n geometrically distinct periodic solutions with minimal period kT .*

Proof Given $n \geq 1$, let p_1, p_2, \dots, p_n be the first n prime numbers. Correspondingly, let $k_{p_1}, k_{p_2}, \dots, k_{p_n}$ be as in Theorem 4. Define

$$k(n) = \max\{k_{p_1}, k_{p_2}, \dots, k_{p_n}\}.$$

By Theorem 4, for every prime number $k \geq k(n)$, equation (6) has periodic solutions $x_{k,p_1}, x_{k,p_2}, \dots, x_{k,p_n}$, with minimal period kT , which make exactly p_1, p_2, \dots, p_n rotations around the origin, respectively, in the period time kT . ■

We immediately deduce the following corollary concerning equation (3).

Corollary 5 *Let $c(t)$ and $e(t)$ be continuous. Assume $e(t)$ has minimal period T , and $\bar{e} < 0$. If $\gamma > 1$ and (8) holds, then the same conclusions of Theorems 4 and 5 hold for system (3).*

References

- [1] D. Bonheure and C. De Coster, Forced singular oscillators and the method of lower and upper solutions, *Topol. Meth. Nonlin. Anal.* 22 (2003), 297–317.
- [2] A. Capietto, J. Mawhin and F. Zanolin, Continuation theorems for periodic perturbations of autonomous systems, *Trans. Amer. Math. Soc.* 329 (1992), 41–72.
- [3] J. Chu, and D. Franco, Non-collision periodic solutions of second order singular dynamical systems, *J. Math. Anal. Appl.* 344 (2008), 898–905.
- [4] V. Coti-Zelati, Dynamical systems with effective-like potential, *Nonlinear Anal.* 12 (1988), 209–222.
- [5] M. del Pino, R. Manasevich and A. Montero, T -periodic solutions for some second order differential equations with singularities, *Proc. Roy. Soc. Edinburgh Sect. A* 120 (1992), 231–243.
- [6] A. Fonda, Periodic solutions for a conservative system of differential equations with a singularity of repulsive type, *Nonlinear Anal.* 24 (1995), 667–676.

- [7] A. Fonda, R. Manasevich and F. Zanolin, Subharmonic solutions for some second-order differential equations with singularities, *SIAM J. Math. Anal.* 24 (1993), 1294–1311.
- [8] A. Fonda and R. Toader, Periodic orbits of radially symmetric Keplerian-like systems: a topological degree approach, *J. Differential Equations* 244 (2008), 3235–3264.
- [9] P. Habets and L. Sanchez, Periodic solutions of dissipative dynamical systems with singular potentials, *Differential Integral Equations* 3 (1990), 1139–1149.
- [10] P. Habets and L. Sanchez, Periodic solutions of some Liénard equations with singularities, *Proc. Amer. Math. Soc.* 109 (1990), 1035–1044.
- [11] A. C. Lazer and S. Solimini, On periodic solutions of nonlinear differential equations with singularities, *Proc. Amer. Math. Soc.* 99 (1987), 109–114.
- [12] J. Mawhin and J. Ward, Periodic solutions of some forced Liénard differential equations at resonance, *Archiv Math.* 41 (1983), 337–351.
- [13] S. Solimini, On forced dynamical systems with a singularity of repulsive type, *Nonlinear Anal.* 14 (1990), 489–500.
- [14] P. J. Torres, Non-collision periodic solutions of forced dynamical systems with weak singularities. *Discrete Contin. Dyn. Syst.* 11 (2004), 693–698.
- [15] E. Zeidler, *Nonlinear Functional Analysis and its Applications*, Vol. 1, Springer, New York–Heidelberg, 1986.

Alessandro Fonda
 Dipartimento di Matematica e Informatica
 Università di Trieste
 P.le Europa 1
 I-34127 Trieste
 Italy
 e-mail: a.fonda@units.it

Rodica Toader
 Dipartimento di Ingegneria Civile e Architettura
 Università di Udine
 Via delle Scienze 208
 I-33100 Udine
 Italy
 e-mail: toader@uniud.it

Mathematics Subject Classification: 34C25

Keywords: periodic solutions, nonlinear dynamics.