

# On a singular periodic Ambrosetti–Prodi problem



Alessandro Fonda<sup>a</sup>, Andrea Sfecci<sup>b,\*</sup>

<sup>a</sup> *Università degli Studi di Trieste, Dipartimento di Matematica e Geoscienze, P.le Europa, 1, I-34127 Trieste, Italy*

<sup>b</sup> *Università Politecnica delle Marche, Dipartimento di Ingegneria Industriale e Scienze Matematiche, Via Breccie Bianche, 12, I-60131 Ancona, Italy*

## ARTICLE INFO

### Article history:

Received 2 August 2016

Accepted 17 October 2016

Communicated by Enzo Mitidieri

### MSC:

34C25

### Keywords:

Periodic solutions

Multiplicity results

Lower and upper solutions

Singularities

Rotating solutions

## ABSTRACT

We investigate the possibility of extending a classical multiplicity result by Fabry, Mawhin and Nkashama to a periodic problem of Ambrosetti–Prodi type having a nonlinearity with possibly one or two singularities. In the second part of the paper we study the existence of periodic rotating solutions for radially symmetric systems with nonlinearities of the same type.

© 2016 Elsevier Ltd. All rights reserved.

## 1. Introduction

In 1972, Ambrosetti and Prodi [1] obtained a multiplicity result for the solutions of a Dirichlet problem associated to an elliptic equation, which can be said to have influenced the research in the field of boundary value problems up to the present days.

Let us recall the result of [1], as refined by Berger and Podolak in [3], by writing the Dirichlet problem as

$$\begin{cases} \Delta u + h(u) = s\varphi_1(x) + w(x) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Here,  $\Omega$  is a bounded domain in  $\mathbb{R}^N$ , while  $\varphi_1(x)$  is the positive eigenfunction associated to the first eigenvalue  $\lambda_1$  of the Laplacian, with Dirichlet boundary conditions, and  $w(x)$  is a suitably smooth function. Assuming  $h : \mathbb{R} \rightarrow \mathbb{R}$  to be twice continuously differentiable and strictly convex, with

$$0 < h'(-\infty) < \lambda_1 < h'(+\infty) < \lambda_2,$$

\* Corresponding author.

E-mail addresses: [a.fonda@units.it](mailto:a.fonda@units.it) (A. Fonda), [sfecci@dipmat.univpm.it](mailto:sfecci@dipmat.univpm.it) (A. Sfecci).

(where  $\lambda_2$  is the second eigenvalue), they proved the existence of an  $s_0 \in \mathbb{R}$  such that

- if  $s < s_0$ , there are no solutions,
- if  $s = s_0$ , there is exactly one solution,
- if  $s > s_0$ , there are exactly two solutions.

Since then, many variants and generalizations have been proposed, see e.g. [2,4,6,15–17,19–21,23–25,28], a far from being exhaustive list. Remarkably, the name *Ambrosetti–Prodi problem* remained attached to all such situations when a multiplicity result structure as the one described above appears.

Searching for an analogue for the periodic problem, Fabry, Mawhin and Nkashama [7] considered in 1986 the second order differential equation

$$x'' + f(x)x' + h(t, x) = s. \tag{E_s}$$

(In this case, the Laplacian is replaced by a second derivative, and the first eigenvalue associated to the periodic problem is equal to zero.) They were able to prove the following Ambrosetti–Prodi type of result.

**Theorem 1.1** (*Fabry–Mawhin–Nkashama*). *Assume  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $h : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  to be continuous functions, with  $T$ -periodicity in the  $t$  variable. If*

$$\lim_{|x| \rightarrow \infty} h(t, x) = +\infty, \quad \text{uniformly in } t \in [0, T],$$

*then there exists an  $s_0 \in \mathbb{R}$  such that*

- *if  $s < s_0$ , there are no  $T$ -periodic solutions,*
- *if  $s = s_0$ , there is at least one  $T$ -periodic solution,*
- *if  $s > s_0$ , there are at least two  $T$ -periodic solutions.*

We will take the above theorem as our starting point, and develop some possible generalizations. In the first part of the paper we focus our attention on the case when the nonlinearities in Eq.  $(E_s)$  are defined only for  $x$  varying in an open interval  $(a, b)$  of  $\mathbb{R}$ , with possibly one or two singularities. Here is our result, extending [Theorem 1.1](#) to such a situation.

**Theorem 1.2.** *Assume  $f : (a, b) \rightarrow \mathbb{R}$  and  $h : \mathbb{R} \times (a, b) \rightarrow \mathbb{R}$  to be continuous functions, with  $T$ -periodicity in the  $t$  variable, such that*

$$\lim_{x \rightarrow a^+} h(t, x) = \lim_{x \rightarrow b^-} h(t, x) = +\infty, \quad \text{uniformly in } t \in [0, T]. \tag{1}$$

*If  $b = +\infty$ , the same conclusion of [Theorem 1.1](#) for Eq.  $(E_s)$  holds. On the other hand, if  $b < +\infty$ , the same is true assuming, in addition, that*

$$f(x) \geq -\eta \quad \text{and} \quad h(t, x) \geq h_m(x), \quad \text{for every } x \in (a, b),$$

*where  $\eta$  is a positive constant and  $h_m : (a, b) \rightarrow \mathbb{R}$  is continuous and such that*

$$\int_c^b h_m(x) dx = +\infty, \tag{2}$$

*for some  $c \in (a, b)$ .*

A few comments on the above statement are in order. Notice that, in the case  $(a, b) = \mathbb{R}$ , [Theorem 1.2](#) reduces to [Theorem 1.1](#). If  $b = +\infty$ , no assumptions besides the continuity are required on the function  $f$ . When  $b < +\infty$ , the repulsive singularity at  $x = b$  has to be sufficiently strong so to ensure that the solutions of  $(E_s)$  cannot collide with it. On the contrary, it is remarkable that the attractive singularity at  $x = a$  does not require an assumption of this type.




Fig. 1. A charged particle  $q^-$  in a capacitor of surface charge density  $\sigma$ , lying between the electric charges  $Q^+$  and  $Q^-$ .

The condition  $f(x) \geq -\eta$  (which could be replaced by the symmetric one  $f(x) \leq \eta$ ) is needed in order to obtain some a priori estimates. However, it would be interesting to know whether it is really necessary.

As a possible example of application of the above theorem, we propose the following physical model describing the dynamics of a charged particle in a periodically varying electric field. We consider a negatively charged particle, freely moving on a straight line between two fixed charged particles, one positive and the other one negative, with  $T$ -periodically varying (not vanishing) charges. We denote by  $q^-$ ,  $Q^+(t)$  and  $Q^-(t)$  the electric charges (in absolute value), respectively. Let  $x = x(t)$  be the position of the freely moving particle, and assume that the fixed charges are placed at  $x = a$  and  $x = b$  respectively, so that  $a < x(t) < b$ , for every  $t$ . We assume that the line of motion is confined between two capacitor plates, as in Fig. 1. The equation of motion is then

$$x'' + k \left( \frac{Q^+(t)}{(x-a)^2} + \frac{Q^-(t)}{(x-b)^2} \right) = s, \quad (3)$$

with  $k = q^-/4\pi\epsilon m$  and  $s = \sigma q^-/m\epsilon$ , where  $m$  is the mass of the free charge,  $\epsilon$  is the dielectric permittivity and  $\sigma$  is the surface charge density of the capacitor. As a consequence of Theorem 1.2, if  $\sigma$  is large enough, Eq. (3) has at least two  $T$ -periodic solutions. A simple physical interpretation of this result can be easily given in the case when the electric charges  $Q^+$  and  $Q^-$  are constant in time: the strong constant force generated by the capacitor balances the attractive force exerted by  $Q^+$ , when the free particle  $x(t)$  is near the position  $x = a$ , and the repulsive force exerted by  $Q^-$ , when  $x(t)$  is near  $x = b$ . Hence there are two equilibria, one near  $x = a$  (unstable) and the other one near  $x = b$  (stable). When the electric charges  $Q^+(t)$  and  $Q^-(t)$  are not constant, but  $T$ -periodic in time, we have a perturbation of the previous situation, if  $\sigma$  is large enough, and the equilibria we have found give rise to the two expected  $T$ -periodic solutions. We recall that the case when  $Q^+$  is replaced by a negative charge has already been considered in [8].

In the second part of the paper, we deal with a system of the type

$$\mathbf{x}'' = (-h(t, |\mathbf{x}|) + s) \frac{\mathbf{x}}{|\mathbf{x}|}. \quad (\mathcal{E}_s)$$

Here  $\mathbf{x} = \mathbf{x}(t) \in \mathbb{R}^N$ , and  $|\cdot|$  denotes the euclidean norm. We will show that the same assumptions considered above on the nonlinearity  $h : \mathbb{R} \times (a, b) \rightarrow \mathbb{R}$ , with  $a \geq 0$ , lead to different types of periodic solutions: some of them will oscillate radially, being those provided by Theorem 1.2. However, new families of periodic solutions will arise, rotating around the origin, completing a revolution in a period time which is a sufficiently large integer multiple of  $T$ .

Let us describe more precisely our result. Writing Eq. ( $\mathcal{E}_s$ ) in polar coordinates, we obtain the system

$$\begin{cases} \rho'' - \frac{\mu^2}{\rho^3} + h(t, \rho) = s, \\ \rho^2 \varphi' = \mu, \end{cases} \quad (\mathcal{R}_s)$$

where  $\mu$  denotes the scalar angular momentum, which is known to be constant along the solutions. This fact is justified by the absence in  $(\mathcal{E}_s)$  of the friction term related to the function  $f$ , which instead was included in the scalar equation  $(E_s)$ . We will only look for solutions with  $\mu > 0$ , since the ones with  $\mu < 0$  can be obtained symmetrically. Notice that the solutions with  $\mu = 0$  (hence with constant  $\varphi$ ) oscillate radially and  $\rho$  solves the scalar equation  $(E_s)$ , with  $f = 0$ . The rotating solutions we are looking for will be such that, for some positive integer  $k$ ,

$$\rho(t + T) = \rho(t), \quad \varphi(t + kT) = \varphi(t) + 2\pi. \tag{4}$$

Notice that such solutions are  $kT$ -periodic, but their radial component is  $T$ -periodic. Let us state our result for the radially symmetric system  $(\mathcal{E}_s)$ .

**Theorem 1.3.** *Let the same assumptions of Theorem 1.2 hold, with  $f = 0$ . Then, there exists an  $s_0 \in \mathbb{R}$  such that, if  $s > s_0$ , system  $(\mathcal{E}_s)$  has two families of rotating periodic solutions with small positive angular momenta. More precisely, there exists a positive integer  $k_s$  such that, for every integer  $k \geq k_s$ , there are two periodic solutions  $(\rho_{k,1}, \varphi_{k,1})$  and  $(\rho_{k,2}, \varphi_{k,2})$  of  $(\mathcal{R}_s)$ , satisfying (4), with positive angular momenta  $\mu_{k,1}$  and  $\mu_{k,2}$ , respectively, such that*

$$\lim_{k \rightarrow \infty} \mu_{k,1} = \lim_{k \rightarrow \infty} \mu_{k,2} = 0.$$

In the case  $b = +\infty$ , if moreover

$$\lim_{r \rightarrow +\infty} \frac{h(t, r)}{r} = 0, \tag{5}$$

then  $(\mathcal{E}_s)$  also admits rotating periodic solutions with large angular momenta. Precisely, there exists a positive integer  $\widehat{k}_s$  such that, for every integer  $k \geq \widehat{k}_s$ , there is a periodic solution  $(\rho_k, \varphi_k)$  of system  $(\mathcal{R}_s)$ , satisfying (4), with positive angular momentum  $\mu_k$ , and with the following properties:

$$\lim_{k \rightarrow \infty} \min \rho_k = +\infty, \quad \lim_{k \rightarrow \infty} \frac{\min \rho_k}{\max \rho_k} = 1, \quad \lim_{k \rightarrow \infty} \mu_k = +\infty.$$

In the above statement, we may have several possible situations: in the case  $b = +\infty$ , if  $a = 0$  we are in the classical case of a Keplerian-type system having only a singularity at the origin. Conversely, if  $a > 0$  we have a *singular sphere*  $\{\rho = a\}$  and the motion takes place outside of it. In the case  $b < +\infty$ , if  $a = 0$  then we have the singularity at the origin and *one singular sphere*  $\{\rho = b\}$  and the motion is confined inside of it. On the other hand, if  $a > 0$ , we have *two singular spheres*,  $\{\rho = a\}$  and  $\{\rho = b\}$ , and the orbits lie in the annular region between them.

Equations of the type  $(\mathcal{E}_s)$  have already been considered in the literature, taking into account several situations. E.g., systems with an attractive singularity of Keplerian type have been studied in [9,12,14]; the case of repulsive singularity has been treated in [10,11,13,26]; bouncing solutions were found in [27]. See also the interesting monograph [29].

The proof of Theorem 1.2 is carried out in Section 2 by the use of lower and upper solutions and topological degree arguments, in the line of the proof given in [7]. Then, in Section 3, we provide the proof of Theorem 1.3, adapting the techniques developed in [10,12]. Finally, in Section 4, we discuss on possible generalizations and extensions of our results.

## 2. Proof of Theorem 1.2

By hypothesis (1), we can define the real number

$$h_0 := \min\{h(t, x) : t \in [0, T], x \in (a, b)\}.$$

If  $s < h_0$ , we cannot have periodic solutions, since otherwise we would have  $x''(t_0) < 0$  at any minimum point  $t_0$ . Let us fix an arbitrary  $\xi \in (a, b)$  and define

$$h_1 := \max\{h(t, \xi) : t \in [0, T]\} \geq h_0.$$

Thanks to assumption (1), for every  $s \in \mathbb{R}$  we can find an interval  $[d_{1,s}, d_{2,s}] \subset (a, b)$  such that

$$h(t, x) > s, \quad \text{for every } (t, x) \in [0, T] \times ((a, d_{1,s}] \cup [d_{2,s}, b)). \quad (6)$$

We can also assume that  $s \mapsto d_{1,s}$  is decreasing and  $s \mapsto d_{2,s}$  is increasing. Notice that  $\alpha \equiv d_{1,s}$  and  $\beta \equiv \xi$  are respectively a lower and an upper solution of  $(E_s)$  with  $\alpha < \beta$ , for every  $s > h_1$ . Then, for every  $s > h_1$  there exists a periodic solution  $x$  of  $(E_s)$  satisfying  $d_{1,s} \leq x(t) \leq \xi$ , for every  $t \in [0, T]$ . Hence, we can define

$$s_0 = \inf\{s \in \mathbb{R} : (E_s) \text{ has a } T\text{-periodic solution}\}. \quad (7)$$

Notice that, by the previous reasoning, we have  $h_0 \leq s_0 \leq h_1$ .

The proof of the following lemma can be traced back to a pioneering paper by Lazer and Solimini [18].

**Lemma 2.1.** *For every  $s > s_0$ , Eq.  $(E_s)$  has a  $T$ -periodic solution.*

**Proof.** We fix  $s > s_0$ . There exists  $\sigma \in [s_0, s)$  such that  $(E_\sigma)$  has a  $T$ -periodic solution, which we denote by  $x_\sigma$ . It is easy to verify that  $x_\sigma$  is an upper solution of  $(E_s)$ . Set  $\alpha_\sigma \in \mathbb{R}$  such that  $a < \alpha_\sigma < d_{1,s}$  and  $\alpha_\sigma < \min x_\sigma$ . Then  $\alpha_\sigma$  is a lower solution of  $(E_s)$ , so that  $(E_s)$  has a  $T$ -periodic solution  $x_s$  satisfying  $\alpha_\sigma \leq x_s(t) \leq x_\sigma(t)$ , for every  $t \in [0, T]$ . The lemma is thus proved. ■

We now prove an a priori estimate for all the possible  $T$ -periodic solutions of  $(E_s)$ , when  $s$  varies in a compact interval.

**Lemma 2.2.** *For every  $\bar{s} > s_0$  there are constants  $\hat{d}_{1,\bar{s}} < \hat{d}_{2,\bar{s}}$  in  $(a, b)$  and  $D_{\bar{s}} > 0$  such that every  $T$ -periodic solution  $x$  of  $(E_s)$  with  $s \in [s_0, \bar{s}]$  must satisfy, for every  $t \in [0, T]$ ,*

$$\hat{d}_{1,\bar{s}} < x(t) < \hat{d}_{2,\bar{s}} \quad \text{and} \quad |x'(t)| < D_{\bar{s}}.$$

**Proof.** Let  $x$  be one such solution. We must have  $\min x > d_{1,\bar{s}}$ , otherwise we would have a negative second derivative at any minimum point. So, we can set  $\hat{d}_{1,\bar{s}} = d_{1,\bar{s}}$ . Let us consider separately the cases  $b = +\infty$  and  $b < +\infty$ .

*Case 1:  $b = +\infty$ .* Integrating Eq.  $(E_s)$  we get

$$\frac{1}{T} \int_0^T h(t, x(t)) dt = s.$$

Introducing the constants  $d_{1,\bar{s}}, d_{2,\bar{s}}$  as in (6), there exists  $t_0 \in [0, T]$  such that  $x(t_0) \in (d_{1,\bar{s}}, d_{2,\bar{s}})$ . Let us denote by  $\bar{x}$  the mean value of  $x$ , i.e.  $\bar{x} = \frac{1}{T} \int_0^T x(t) dt$ , so that  $\tilde{x}(t) = x(t) - \bar{x}$  has zero mean. Multiplying  $(E_s)$  by  $\tilde{x}$  we get

$$\begin{aligned} \|x'\|_2^2 &= \int_0^T \tilde{x}(t) h(t, x(t)) dt = \int_0^T \tilde{x}(t) (h(t, x(t)) - h_0) dt \\ &\leq \|\tilde{x}\|_\infty \int_0^T (h(t, x(t)) - h_0) dt \leq \|\tilde{x}\|_\infty T(\bar{s} - h_0). \end{aligned}$$

Let  $t_1 \in [0, T]$  be such that  $\tilde{x}(t_1) = 0$ . Then, for every  $t \in [t_1, t_1 + T]$ ,

$$|\tilde{x}(t)| \leq \left| \int_{t_1}^t \tilde{x}'(\tau) d\tau \right| \leq \int_0^T |x'(\tau)| d\tau \leq \sqrt{T} \|x'\|_2.$$

By the previous estimates, we get  $\|x'\|_2 \leq T^{3/2}(\bar{s} - h_0)$ , so that, for every  $t \in [0, T]$ ,

$$x(t) = x(t_0) + \int_{t_0}^t x'(\tau) d\tau \leq x(t_0) + \sqrt{T}\|x'\|_2 < d_{2,\bar{s}} + T^2(\bar{s} - h_0).$$

Hence, setting  $\widehat{d}_{2,\bar{s}} = d_{2,\bar{s}} + T^2(\bar{s} - h_0)$ , we have  $x(t) < \widehat{d}_{2,\bar{s}}$ , for every  $t \in [0, T]$ .

*Case 2:*  $b < +\infty$ . We introduce the energy  $E(x, y) = y^2/2 + H(x)$ , where  $H(x) = \int_c^x (h_m(v) - \bar{s}) dv$ . There exists a constant  $H_0$  such that  $H(v) \leq H_0$  for every  $v \in [d_{1,\bar{s}}, d_{2,\bar{s}}]$ . By (2), it is possible to find a  $\widehat{d}_{2,\bar{s}} \in (d_{2,\bar{s}}, b)$  such that  $H(v) > H_0 e^{2\eta T}$ , for every  $v \in [\widehat{d}_{2,\bar{s}}, b)$ .

Assume  $\max x > d_{2,\bar{s}}$ . Then, by (6), there exist  $t_1 < t_2$  such that  $x'(t_1) = x'(t_2) = 0$ ,  $x(t_2) = \max x$  and  $x'(t) > 0$  for every  $t \in (t_1, t_2)$ , and  $x(t_1) \in [d_{1,\bar{s}}, d_{2,\bar{s}}]$ . A computation gives, for every  $t \in (t_1, t_2)$ ,

$$\begin{aligned} \frac{d}{dt} E(x(t), x'(t)) &= -x'(t)[f(x(t))x'(t) + h(t, x(t)) - s - h_m(x(t)) + \bar{s}] \\ &\leq \eta (x'(t))^2 \leq 2\eta E(x(t), x'(t)). \end{aligned}$$

Hence,

$$H(x(t_2)) = E(x(t_2), x'(t_2)) \leq E(x(t_1), x'(t_1)) e^{2\eta(t_2-t_1)} \leq H_0 e^{2\eta T},$$

thus giving us  $x(t_2) < \widehat{d}_{2,\bar{s}}$ .

The proof of the derivative estimate follows easily from the validity of a Nagumo condition. Indeed, by the previously proved estimates, we get the existence of a positive constant  $C$  such that, for every  $T$ -periodic solution  $x$  of  $(E_s)$ , one has

$$x''(t) \leq C(|x'(t)| + 1), \quad \text{for every } t \in [0, T].$$

The application of Gronwall Lemma ends the proof. ■

We remark that it is possible to assume, without loss of generality, that  $s \mapsto \widehat{d}_{1,s}$  is decreasing,  $s \mapsto \widehat{d}_{2,s}$  and  $s \mapsto D_s$  are increasing.

We now define our functional setting. Let  $X = C([0, T])$  be the set of continuous functions, and let  $\mathcal{L} : \mathcal{D}(\mathcal{L}) \rightarrow X$  be the operator defined as

$$\begin{aligned} \mathcal{D}(\mathcal{L}) &= \{x \in C^2([0, T]) : x(0) = x(T), x'(0) = x'(T)\}, \\ \mathcal{L}x &= x'' - x. \end{aligned}$$

Setting  $Y = C^1([0, T])$ , we define on

$$Y_{(a,b)} = \{x \in Y : a < x(t) < b, \text{ for every } t \in [0, T]\}$$

the Nemytskii operator  $N_s : Y_{(a,b)} \rightarrow X$  as

$$(N_s x)(t) = -f(x(t))x'(t) - h(t, x(t)) + s - x(t).$$

We thus have that  $x$  is a  $T$ -periodic solution of  $(E_s)$  if and only if it solves the equation

$$\mathcal{L}x = N_s x,$$

with  $x \in \mathcal{D}(\mathcal{L}) \cap Y_{(a,b)}$ . Fix  $\bar{s} > s_0$  and define the set

$$\overline{\Xi}_{\bar{s}} = \left\{ x \in Y : \widehat{d}_{1,\bar{s}} < x(t) < \widehat{d}_{2,\bar{s}} \text{ and } |x'(t)| < D_{\bar{s}}, \text{ for every } t \in [0, T] \right\}.$$

By standard arguments, for every  $s < \bar{s}$ , the function  $\Psi_s = \mathcal{L}^{-1} \circ N_s : \overline{\Xi}_{\bar{s}} \rightarrow Y$  is a completely continuous operator, and its fixed points are the  $T$ -periodic solutions of  $(E_s)$ . By Lemma 2.2, we have

that  $0 \notin (I - \Psi_s)(\partial \Xi_{\bar{s}})$ , so that the degree of  $I - \Psi_s$  on the open and bounded set  $\Xi_{\bar{s}} \subset Y$  is well defined, for every  $s \leq \bar{s}$ . Recalling that for every  $s < s_0$  there are no  $T$ -periodic solutions of  $(E_s)$ , using the homotopy invariance property of the degree, we have

$$d(I - \Psi_s, \Xi_{\bar{s}}) = 0, \quad \text{for every } s \leq \bar{s}.$$

Now, for every  $\varepsilon > 0$ , consider a  $T$ -periodic solution  $\beta_\varepsilon$  of equation  $(E_{s_0+\varepsilon})$ , whose existence is guaranteed by Lemma 2.1. If  $s_0 + \varepsilon < s \leq \bar{s}$ , then  $\alpha_\varepsilon \equiv \widehat{a}_{1,\bar{s}}$  and  $\beta_\varepsilon$  are respectively a lower and an upper solution of  $(E_s)$ , and  $\alpha_\varepsilon < \beta_\varepsilon(t)$  for every  $t \in [0, T]$ , by Lemma 2.2. Set

$$\Omega_\varepsilon^1 = \{x \in Y : \alpha_\varepsilon < x(t) < \beta_\varepsilon(t) \text{ and } |x'(t)| < D_{\bar{s}}, \text{ for every } t \in [0, T]\},$$

a subset of  $\Xi_{\bar{s}}$ , by Lemma 2.2. We now prove that there are no  $T$ -periodic solutions of  $(E_s)$  belonging to  $\partial \Omega_\varepsilon^1$ , if  $s \in (s_0 + \varepsilon, \bar{s}]$ . Let  $x$  be a  $T$ -periodic solution of  $(E_s)$  such that  $\alpha_\varepsilon \leq x(t) \leq \beta_\varepsilon(t)$ , for every  $t \in [0, T]$ . Arguing as above, we see that such a solution cannot have  $\alpha_\varepsilon$  as a minimum. Conversely, suppose that there exists a  $\tau \in [0, T]$  such that  $x(\tau) - \beta_\varepsilon(\tau) = 0$ . Then,  $\tau$  is a point of maximum for  $x(t) - \beta_\varepsilon(t)$ , hence  $x'(\tau) - \beta'_\varepsilon(\tau) = 0$ , and we have

$$x''(\tau) - \beta''_\varepsilon(\tau) = -f(x(\tau))x'(\tau) - h(\tau, x(\tau)) + s + f(\beta_\varepsilon(\tau))\beta'_\varepsilon(\tau) + h(\tau, \beta_\varepsilon(\tau)) - s_0 - \varepsilon = s - s_0 - \varepsilon > 0, \quad (8)$$

leading to a contradiction. Hence, by a standard result in lower and upper solution theory (see, e.g., [5]),

$$d(I - \Psi_s, \Omega_\varepsilon^1) = 1, \quad \text{for every } s \in (s_0 + \varepsilon, \bar{s}]. \quad (9)$$

We now define  $\Omega_\varepsilon^2 = \Xi_{\bar{s}} \setminus \overline{\Omega_\varepsilon^1}$ . By the additivity property of the degree,

$$d(I - \Psi_s, \Omega_\varepsilon^2) = -1, \quad \text{for every } s \in (s_0 + \varepsilon, \bar{s}]. \quad (10)$$

Hence, since the choice of  $\varepsilon$  is arbitrary, for every  $s \in (s_0, \bar{s}]$  there are at least two  $T$ -periodic solutions of  $(E_s)$ , one in  $\Omega_\varepsilon^1$  and the second one in  $\Omega_\varepsilon^2$ , simply choosing  $\varepsilon < s - s_0$ . Since we can consider  $\bar{s} > s_0$  arbitrarily large, we have thus proved that there exist two  $T$ -periodic solutions of  $(E_s)$ , for every  $s > s_0$ .

**Remark 2.3.** We have proved that there are at least two  $T$ -periodic solutions, but there could be many more: we will discuss this issue in Section 4. Let us mention here that, among all the  $T$ -periodic solutions, it is always possible to find two of them which are ordered. Indeed, let  $x_1$  and  $x_2$  be the two  $T$ -periodic solutions found above. Then the function  $\bar{\beta} = \min\{x_1, x_2\}$  is an upper solution of  $(E_s)$ , and therefore there exists a  $T$ -periodic solution  $\tilde{x}_1$  between  $\alpha_\varepsilon$  and  $\bar{\beta}$ , cf. [5, Chapter VI].

In order to prove the existence of at least one periodic solution of  $(E_{s_0})$ , we consider a strictly decreasing sequence  $(s_n)_n$  with  $\lim_n s_n = s_0$ . For every  $n$ , let  $x_n$  be a solution of  $(E_{s_n})$ . By Lemma 2.2, we have that  $(x_n)_n$  is contained in  $\Xi_{\bar{s}}$ . Moreover, by the fact that  $x_n$  solves the differential equation  $(E_{s_n})$ , we have that  $(x_n)_n$  is bounded in  $C^2([0, T])$ , so that, by the Ascoli–Arzelà Theorem, it  $C^1$ -converges up to subsequences to some  $x \in \overline{\Xi_{\bar{s}}}$ . Since  $x_n = \Psi_{s_n}(x_n)$ , passing to the limit, we obtain  $x = \Psi_{s_0}(x)$ , so that  $x$  solves  $(E_{s_0})$ . The proof of Theorem 1.2 is thus completed.

### 3. Proof of Theorem 1.3

Setting  $X = C([0, T])$  and

$$X_{(a,b)} = \{\rho \in X : a < \rho(t) < b, \text{ for every } t \in [0, T]\},$$

the Nemytskii operator  $N_{s,\mu} : X_{(a,b)} \rightarrow X$  can now be defined as

$$(N_{s,\mu} \rho)(t) := \frac{\mu^2}{\rho^3(t)} - h(t, \rho(t)) + s - \rho(t).$$

Let  $\Omega$  be an open bounded subset of  $X_{(a,b)}$  such that  $\bar{\Omega} \subset X_{(a,b)}$ . The operator  $\Psi_{s,\mu} = \mathcal{L}^{-1} \circ N_{s,\mu} : \bar{\Omega} \rightarrow X$  is completely continuous and its fixed points correspond to  $T$ -periodic solutions of the first equation in  $(\mathcal{R}_s)$ .

The following theorem is a variant of [12, Theorem 2].

**Theorem 3.1.** *Assume that there are no fixed points of  $\Psi_{s,0}$  on  $\partial\Omega$ , and that  $d(I - \Psi_{s,0}, \Omega) \neq 0$ . Then, there exists a  $\bar{k} \geq 1$  such that, for every integer  $k \geq \bar{k}$ , system  $(\mathcal{R}_s)$  has a  $kT$ -periodic solution  $(\rho_k, \varphi_k)$  satisfying (4). Moreover,  $\rho_k$  belongs to  $\Omega$  and, if  $\mu_k$  denotes the associated angular momentum, then*

$$\lim_{k \rightarrow \infty} \mu_k = 0.$$

The proof of Theorem 3.1 is completely analogous of the one provided in [12]. It can also be carried out by suitably modifying the nonlinearity  $h$ , in the following way. Take an interval  $[c, d] \subset (a, b)$  such that  $\bar{\Omega} \subset X_{(c,d)}$ , and a function  $\tilde{h} : \mathbb{R} \times (0, +\infty) \rightarrow \mathbb{R}$  such that  $\tilde{h} = h$  on  $\mathbb{R} \times (c, d)$ . Replacing  $h$  with  $\tilde{h}$  in  $(\mathcal{R}_s)$  we are brought back to the setting of a Newtonian system already considered in [12, Theorem 2], and the result follows.

Going back to the first equation in  $(\mathcal{R}_s)$ , we first study the situation when  $\mu = 0$ . Following the proof of the first part of Theorem 1.3, we fix  $\bar{s} > s_0$  and define

$$\Xi_{\bar{s}} = \{ \rho \in X : \hat{d}_{1,\bar{s}} < \rho(t) < \hat{d}_{2,\bar{s}}, \text{ for every } t \in [0, T] \}.$$

Taking  $s \in (s_0, \bar{s}]$ , we can choose  $\varepsilon < \bar{s} - s_0$  and define

$$\Omega_\varepsilon^1 = \{ \rho \in X : \alpha_\varepsilon < \rho(t) < \beta_\varepsilon(t), \text{ for every } t \in [0, T] \},$$

a subset of  $\Xi_{\bar{s}}$ , where we have used the notation of the previous section for  $\alpha_\varepsilon$  and  $\beta_\varepsilon(t)$ . Finally, we set  $\Omega_\varepsilon^2 = \Xi_{\bar{s}} \setminus \bar{\Omega}_\varepsilon^1$ . We thus obtain the analogues of formulas (9) and (10), i.e.,

$$d(I - \Psi_{s,0}, \Omega_\varepsilon^1) = 1, \quad d(I - \Psi_{s,0}, \Omega_\varepsilon^2) = -1,$$

for every  $s \in (s_0 + \varepsilon, \bar{s}]$ , and we can apply Theorem 3.1 with  $\Omega = \Omega_\varepsilon^1$  and  $\Omega = \Omega_\varepsilon^2$ , thus finding the two required families of periodic solutions with a small angular momentum.

When  $b = +\infty$ , the proof of the second part of the statement follows directly from [10, Theorem 1.2], after suitably modifying the function  $h : \mathbb{R} \times (a, +\infty) \rightarrow \mathbb{R}$  to some function  $\hat{h} : \mathbb{R} \times (0, +\infty) \rightarrow \mathbb{R}$  such that  $\hat{h} = h$  on  $\mathbb{R} \times (c, +\infty)$ , for some  $c > a$ .

#### 4. Final remarks

As observed in [5], the continuity assumption on the function  $h(t, x)$  can be replaced by  $L^1$ -Carathéodory conditions, provided an extra hypothesis is fulfilled (cf. assumption (A) in [5, Theorem VI-1.2 and Corollary VI-1.3]).

Concerning the multiplicity of solutions of Eq.  $(E_s)$ , their exact number can be established, in the case  $b = +\infty$ , if  $f = 0$  and  $h(t, \cdot)$  is strictly convex, with

$$\frac{h(t, x_1) - h(t, x_2)}{x_1 - x_2} < \left( \frac{2\pi}{T} \right)^2, \quad \text{for every } x_1 \neq x_2 \text{ and } t \in [0, T].$$

The proof can be carried out applying [5, Theorem VI-1.4], after suitably modifying and extending  $h(t, \cdot)$  on the interval  $(-\infty, a + \varepsilon)$ , for some  $\varepsilon > 0$ , so to obtain a strictly convex function defined on the whole real



line. Notice that a larger number of solutions of Eq.  $(E_s)$  can arise when the nonlinearity  $h$  “oscillates”: for example, if  $h(t, x) = h(x)$ , the differential equation  $x'' + f(x)x' + h(x) = s$  may clearly have several constant solutions.

Let us argue on the stability of the two  $T$ -periodic solutions found in [Theorem 1.2](#), when  $b = +\infty$ . For an equation of the type

$$x'' + cx' + h(x) = s + p(t),$$

with  $c > 0$ , assume  $h$  to be twice continuously differentiable and strictly convex, with  $h'(+\infty) \leq (\pi/T)^2 + c^2/4$ . By a truncation argument, it is possible to reduce to [\[22, Theorem 2.1\]](#) showing that, also in this case, there are exactly two  $T$ -periodic solutions for  $s$  large, one of which is asymptotically stable, and the other one unstable (see also [\[23\]](#) for further considerations). We omit the details, for brevity.

As a further remark, we notice that, adapting the proof of [Theorem 1.2](#), it is easy to see that the following result also holds.

**Corollary 4.1.** *Assume*

$$\lim_{x \rightarrow a^+} h(t, x) = +\infty, \quad \text{uniformly in } t \in [0, T].$$

*Then, there exists at least one  $T$ -periodic solution of Eq.  $(E_s)$ , provided that  $s$  is sufficiently large.*

Indeed, in such a situation it is possible to find a lower and an upper solution of  $(E_s)$ , for large values of  $s$ . We emphasize that, being now  $h$  not necessarily bounded from below, the value  $s_0$  introduced in [\(7\)](#) can be equal to  $-\infty$ . As an example of application, a physical model similar to the one described in [Fig. 1](#) can be considered, by dropping the charge  $Q^-$ , or replacing it with a positive one. In the former case  $s_0 \in \mathbb{R}$ , and in the latter  $s_0 = -\infty$ .

Some refinements of [Theorem 1.3](#) can also be obtained: for instance, when the exact number of solutions of [Theorem 1.2](#) is known, we can preserve an order structure of the rotating solutions provided there. More precisely, referring to [Remark 2.3](#), the two solutions given by [Theorem 1.2](#), for  $s > s_0$ , can be assumed to be ordered as  $x_1 \prec x_2$ , meaning that  $x_1(t) \leq x_2(t)$  for every  $t \in [0, T]$ , and  $x_1 \neq x_2$ . Defining the average function  $r = (x_1 + x_2)/2$ , we have that  $x_1 \prec r \prec x_2$ . Using the excision property of the degree, we can replace, in the proof of [Theorem 1.3](#), the sets  $\Omega_\varepsilon^1$  and  $\Omega_\varepsilon^2$ , respectively, by

$$\Gamma^1 = \{\rho \in \Xi_s : \rho \prec r\} \quad \text{and} \quad \Gamma^2 = \{\rho \in \Xi_s : r \prec \rho\}.$$

In this way, for every couple of solutions  $(\rho_{k,1}, \varphi_{k,1})$  and  $(\rho_{j,2}, \varphi_{j,2})$  emerging from  $\Gamma^1$  and  $\Gamma^2$ , respectively, we have that  $\rho_{k,1} \prec \rho_{j,2}$ .

Finally, under the assumptions of [Corollary 4.1](#), arguing as in the proof of [Theorem 1.3](#), it is possible to find at least one family of rotating solutions for the radially symmetric system  $(\mathcal{E}_s)$ , with a small angular momentum.

## Acknowledgements

The authors are grateful to the referee for the careful reading of the paper and for useful suggestions. This work was partially supported by the INdAM-GNAMPA project (2016) *Problemi differenziali non lineari: esistenza, molteplicità e proprietà qualitative delle soluzioni*.

## References

- [1] A. Ambrosetti, G. Prodi, On the inversion of some differentiable mappings with singularities between Banach spaces, *Ann. Mat. Pura Appl.* 93 (1972) 231–246.
- [2] C. Bereanu, An Ambrosetti–Prodi–type result for periodic solutions of the telegraph equation, *Proc. Roy. Soc. Edinburgh Sect. A* 138 (2008) 719–724.

- [3] M.S. Berger, E. Podolak, On the solutions of a nonlinear Dirichlet problem, *Indiana Univ. Math. J.* 24 (1975) 837–846.
- [4] J. Berkovits, Ambrosetti–Prodi type multiplicity result for a wave equation with sublinear nonlinearity, *J. Math. Anal. Appl.* 332 (2007) 691–699.
- [5] C. De Coster, P. Habets, *Two-Point Boundary Value Problems: Lower and Upper Solutions*, Elsevier, Amsterdam, 2006.
- [6] D.G. de Figueiredo, On the superlinear Ambrosetti–Prodi problem, *Nonlinear Anal.* 8 (1984) 655–665.
- [7] C. Fabry, J. Mawhin, M.N. Nkashama, A multiplicity result for periodic solutions of forced nonlinear second order ordinary differential equations, *Bull. Lond. Math. Soc.* 18 (1986) 173–180.
- [8] A. Fonda, R. Manásevich, F. Zanolin, Subharmonic solutions for some second-order differential equations with singularities, *SIAM J. Math. Anal.* 24 (1993) 1294–1311.
- [9] A. Fonda, R. Toader, Periodic orbits of radially symmetric Keplerian-like systems: A topological degree approach, *J. Differential Equations* 244 (2008) 3235–3264.
- [10] A. Fonda, R. Toader, Periodic orbits of radially symmetric systems with a singularity: the repulsive case, *Adv. Nonlinear Stud.* 11 (2011) 853–874.
- [11] A. Fonda, R. Toader, Radially symmetric systems with a singularity and asymptotically linear growth, *Nonlinear Anal.* 74 (2011) 2485–2496.
- [12] A. Fonda, R. Toader, Periodic solutions of radially symmetric perturbations of Newtonian systems, *Proc. Amer. Math. Soc.* 140 (2012) 1331–1341.
- [13] A. Fonda, R. Toader, F. Zanolin, Periodic solutions of singular radially symmetric systems with superlinear growth, *Ann. Mat. Pura Appl.* 191 (2012) 181–204.
- [14] A. Fonda, A.J. Ureña, Periodic, subharmonic, and quasi-periodic oscillations under the action of a central force, *Discrete Contin. Dyn. Syst.* 29 (2011) 169–192.
- [15] S. Fučík, Remarks on a result by A. Ambrosetti and G. Prodi, *Boll. Unione Mat. Ital.* (4) 11 (1975) 259–267.
- [16] P. Hess, On a nonlinear elliptic boundary value problem of the Ambrosetti–Prodi type, *Boll. Unione Mat. Ital. A* (5) 17 (1980) 187–192.
- [17] J.L. Kazdan, F.W. Warner, Remarks on some quasilinear elliptic equations, *Comm. Pure Appl. Math.* 28 (1975) 567–597.
- [18] A.C. Lazer, S. Solimini, On periodic solutions of nonlinear differential equations with singularities, *Proc. Amer. Math. Soc.* 88 (1987) 109–114.
- [19] J. Mawhin, Ambrosetti–Prodi type results in nonlinear boundary value problems, in: *Differential Equations and Mathematical Physics* (Birmingham, 1986), in: *Lect. Notes Math.*, vol. 1285, Springer, Berlin, 1987, pp. 280–313.
- [20] J. Mawhin, The periodic Ambrosetti–Prodi problem for nonlinear perturbations of the  $p$ -Laplacian, *J. Eur. Math. Soc.* 8 (2006) 375–388.
- [21] F. Obersnel, P. Omari, On the Ambrosetti–Prodi problem for first order scalar periodic ODEs, in: *Applied and Industrial Mathematics in Italy*, in: *Ser. Adv. Math. Appl. Sci.*, vol. 69, World Sci. Publ, Hackensack, NJ, 2005, pp. 404–415.
- [22] R. Ortega, Stability and index of periodic solutions of an equation of Duffing type, *Boll. Unione Mat. Ital. B* (7) 3 (1989) 533–546.
- [23] R. Ortega, Stability of a periodic problem of Ambrosetti–Prodi type, *Differential Integral Equations* 3 (1990) 275–284.
- [24] R. Ortega, M. Tarallo, Almost periodic equations and conditions of Ambrosetti–Prodi type, *Math. Proc. Cambridge Philos. Soc.* 135 (2003) 239–254.
- [25] C. Rebelo, F. Zanolin, Multiplicity results for periodic solutions of second order ODEs with asymmetric nonlinearities, *Trans. Amer. Math. Soc.* 348 (1996) 2349–2389.
- [26] A. Sfecci, Double resonance for one-sided superlinear or singular nonlinearities, *Ann. Mat. Pura Appl.* 195 (2016) 2007–2025.
- [27] A. Sfecci, Periodic impact motions at resonance of a particle bouncing on spheres and cylinders, preprint arXiv:1504.03457.
- [28] E. Sovrano, F. Zanolin, The Ambrosetti–Prodi periodic problem: different routes to complex dynamics, preprint, 2016.
- [29] P.J. Torres, *Mathematical Models with Singularities. A Zoo of Singular Creatures*, Atlantis Press, Paris, 2015.