



# A Lower/Upper Solutions Result for Generalised Radial $p$ -Laplacian Boundary Value Problems

Alessandro Fonda , Natnael Gezahegn Mamo ,  
Franco Obersnel  and Andrea Sfecci 

**Abstract.** We provide existence results to some planar nonlinear boundary value problems, in the presence of lower and upper solutions. Our results apply to a class of systems generalising radial elliptic equations driven by the  $p$ -Laplace operator, and to some problems involving the Laplace–Beltrami operator on the sphere. After extending the definition of lower and upper solutions to the planar system, we prove our results by a shooting method involving a careful analysis of the solutions in the phase plane.

**Mathematics Subject Classification.** 34B15.

**Keywords.** Radial elliptic equations, lower and upper solutions, shooting method,  $p$ -Laplacian, Laplace–Beltrami operator.

## 1. Introduction and Main Results

The lower and upper solutions method is a classical tool for studying boundary value problems associated with ordinary and partial differential equations of different types. Since the pioneering works by Picard [11], Scorza Dragoni [13], Nagumo [10], thousands of papers have employed it to study existence, multiplicity, localisation and stability properties of the solutions of first- and second-order problems. See, e.g. [4] for a classical monograph on the topic. To present a simple but illustrative example, let us consider the Neumann problem

$$x'' = g(t, x), \quad x'(0) = x'(1) = 0. \quad (1)$$

---

Natnael Gezahegn Mamo expresses his gratitude to the Department of Mathematics, Addis Ababa University and International Science Program (ISP), Uppsala University, for their financial support for the research visit at the Department of Mathematics and Geosciences of the University of Trieste. In addition, he would like to thank the Department for graciously hosting him from May to August 2022.

For this problem, a lower solution  $\alpha : [0, 1] \rightarrow \mathbb{R}$  is defined as a  $C^2$ -function satisfying  $\alpha''(t) \geq g(t, \alpha(t))$ , for every  $t \in [0, 1]$ , and  $\alpha'(0) \geq 0 \geq \alpha'(1)$ . An upper solution  $\beta$  is similarly defined by reversing the inequalities. If we set  $y = x'$ , problem (1) is equivalent to the planar system

$$x' = f(t, y), \quad y' = g(t, x), \quad y(0) = y(1) = 0, \tag{2}$$

where  $f(t, y) = y$ . The relations defining  $\alpha$  and  $\beta$  translate into

$$y'_\alpha(t) \geq g(t, \alpha(t)), \quad \text{for every } t \in [0, 1], \quad y_\alpha(0) \geq 0 \geq y_\alpha(1),$$

where  $y_\alpha = \alpha'$ ,

$$y'_\beta(t) \leq g(t, \beta(t)), \quad \text{for every } t \in [0, 1], \quad y_\beta(0) \leq 0 \leq y_\beta(1),$$

where  $y_\beta = \beta'$ .

With similar models in mind, Fonda and Toader in [8] have extended to planar systems the definitions of lower and upper solutions for a wide class of problems and for general equations of the form

$$x' = f(t, x, y), \quad y' = g(t, x, y).$$

In particular, we refer to [8, §2] for the definition in the periodic setting (see also [6, §2] for the non-smooth case), and to [7, §2] for separated boundary value problems (see also [5, §4]).

In [7] (see, e.g. Theorems 5 and Corollary 10), assuming the existence of a lower solution  $\alpha$  and an upper solution  $\beta$ , with  $\alpha \leq \beta$ , it is proved that there exists a solution  $(x, y)$  of (2) satisfying  $\alpha \leq x \leq \beta$ . In this paper, we are interested in extending this result to systems of the type

$$x' = f(t, y), \quad (a(t)y)' = g(t, x), \tag{3}$$

motivated by the study of radial weighted  $p$ -Laplacian differential equations as, e.g. in [1–3, 9]. Consider, without loss of generality, the equation in the unitary ball  $\mathcal{B}$

$$\operatorname{div}(\eta(|x|)|\nabla v|^{p-2}\nabla v) = h(|x|, v), \tag{4}$$

where  $\eta : [0, 1] \rightarrow \mathbb{R}^+$  is a strictly positive smooth radial weight,  $h : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous, and  $p > 1$ . We are interested in finding radial solutions of (4) of the form  $v(x) = u(|x|) = u(r)$ . The function  $u : [0, 1] \rightarrow \mathbb{R}$  satisfies the equation

$$(a(r)|u'|^{p-2}u')' = g(r, u), \quad r \in ]0, 1], \tag{5}$$

with  $a(r) = r^{N-1}\eta(r)$  and  $g(r, u) = r^{N-1}h(r, u)$ , it is continuously differentiable with  $u'(0) = 0$ , and  $a(\cdot)|u'|^{p-2}u'$  is also continuously differentiable.

Denoting by  $q > 1$  the conjugate exponent of  $p$ , satisfying  $\frac{1}{p} + \frac{1}{q} = 1$ , Eq. (5) is equivalent to the system

$$u' = |y|^{q-2}y, \quad (a(r)y)' = g(r, u),$$

which is a special form of (3). Note that the function  $a(r)$  vanishes at  $r = 0$ , creating a singularity for our problem, and this fact generates a main difficulty in our study. The problem of the presence of the singularity, concerning existence, uniqueness and continuous dependence on initial data, for the Cauchy problems associated with the second-order differential equation (5), was already faced in [9] (in the Appendix), see also [2–4]. In the appendix of this

paper, we present the corresponding discussion for system (3). Moreover, we can also allow the possibility of having a second singularity at  $r = 1$ .

We consider the mixed boundary conditions problem

$$\begin{cases} x' = f(t, y), & (a(t)y)' = g(t, x), \\ y(0) = 0 = x(1) \sin \theta + y(1) \cos \theta, \end{cases} \tag{6}$$

with  $\theta \in ]-\frac{\pi}{2}, \frac{\pi}{2}]$ . Having in mind the radial problem, it is natural to assume the Neumann condition  $y(0) = 0$  at the left endpoint of our interval. Concerning the right endpoint, notice that, in case  $\theta = 0$ , we have a Neumann-type boundary condition, while in case  $\theta = \frac{\pi}{2}$ , we are dealing with a Dirichlet-type condition.

Let  $a : [0, 1] \rightarrow \mathbb{R}$  satisfy the following assumptions:

- (A1)  $a \in C^1([0, 1])$ ;
- (A2)  $a(t) > 0$ , for all  $t \in ]0, 1[$ ;
- (A3)  $a(0) = 0$ , and there exists  $\rho_0 \in ]0, 1[$  such that

$$a'(t) \geq 0, \quad \text{for every } t \in [0, \rho_0].$$

*Remark 1.* Assume  $N \geq 2$  and  $\eta : [0, 1] \rightarrow \mathbb{R}^+$  is strictly positive and continuously differentiable on  $[0, 1]$ . Then, the function  $a(r) = r^{N-1}\eta(r)$  introduced in (5) satisfies assumptions (A1), (A2) and (A3).

We now state our first result, where the existence of a pair of well-ordered lower and upper solutions is assumed, referring to Sect. 2.1 for their definitions.

**Theorem 2.** *Assume  $f, g : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous, locally Lipschitz continuous in the second variable, and  $a : [0, 1] \rightarrow \mathbb{R}$  satisfies (A1), (A2) and (A3). Assume that the function  $\hat{g} : ]0, 1[ \times \mathbb{R} \rightarrow \mathbb{R}$ , defined by*

$$\hat{g}(t, x) = \frac{1}{a(t)}g(t, x), \tag{7}$$

*can be continuously extended to  $[0, 1] \times \mathbb{R}$ . Suppose further that there exist a lower solution  $\alpha$  and an upper solution  $\beta$  for problem (6), satisfying  $\alpha \leq \beta$ . Then, problem (6) has a solution  $(x, y)$  such that  $\alpha \leq x \leq \beta$ .*

Concerning the proof of the above theorem, we present an alternative approach to the standard application of degree theory, based on a shooting method, after a careful phase plane analysis of the solutions.

*Remark 3.* We underline that if we replace assumptions (A2) and (A3) with the hypothesis  $a(t) > 0$ , for all  $t \in [0, 1]$ , then the conclusion of Theorem 2 can be proved with simpler computations.

If we are only interested in the Neumann problem (6), with  $\theta = 0$ , we can weaken the assumptions on the function  $a$  by allowing  $a(1) = 0$ . We shall assume that  $a : [0, 1] \rightarrow \mathbb{R}$  satisfies (A1) and

- (A2)'  $a(t) > 0$ , for all  $t \in ]0, 1[$ ;
- (A3)'  $a(0) = 0, a(1) = 0$ , and there exist  $\rho_0 \leq \rho_1$  in  $]0, 1[$  such that

$$a'(t) \geq 0 \text{ for every } t \in [0, \rho_0] \text{ and } a'(t) \leq 0 \text{ for every } t \in [\rho_1, 1].$$

An example of a function  $a$  satisfying these assumptions is

$$a(t) = \sin^{N-2}(\pi t), \quad N \geq 3. \tag{8}$$

It arises, e.g. when dealing with the Laplace–Beltrami operator on the sphere  $\mathbb{S}^{N-1} \subseteq \mathbb{R}^N$ , if we are looking for solutions depending only on the latitude  $\varphi = \pi t$  (asking for symmetry with respect to all the other angle variables). In this case, the problem we need to solve is the following

$$\begin{cases} (\sin^{N-2}(\pi t)x')' = \sin^{N-2}(\pi t)g(t, x), \\ x'(0) = 0 = x'(1), \end{cases} \tag{9}$$

which is a special form of (6), with  $\theta = 0$ , the function  $a$  defined by (8) and the function  $g$  replaced by  $\sin^{N-2}(\pi t)g(t, x)$ .

Concerning this new setting, we can state our second result.

**Theorem 4.** *Assume  $f, g : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  are continuous, locally Lipschitz continuous in the second variable, and  $a : [0, 1] \rightarrow \mathbb{R}$  satisfies (A1), (A2)' and (A3)'. Assume that the function  $\hat{g} : ]0, 1[ \times \mathbb{R} \rightarrow \mathbb{R}$ , defined by (7) can be continuously extended to  $[0, 1] \times \mathbb{R}$ . Suppose further that there exist a lower solution  $\alpha$  and an upper solution  $\beta$  for problem (6), with  $\theta = 0$ , satisfying  $\alpha \leq \beta$ . Then, problem (6) with  $\theta = 0$  has a solution  $(x, y)$  such that  $\alpha \leq x \leq \beta$ .*

The proof of this second theorem will be provided through a *double shooting method*.

*Remark 5.* If the functions  $f$  and  $g$  are only continuous, similar results can still be proved, adapting the approximation technique used in [8]. However, in this case, we need to assume the existence of strict lower and upper solutions  $\alpha$  and  $\beta$  satisfying  $\alpha(t) < \beta(t)$  for all  $t \in ]0, 1[$ .

## 2. Preliminaries

### 2.1. Lower and Upper Solutions

We now provide the definitions of lower and upper solutions for problem (6), thus extending the ones given in [5–7].

**Definition 6.** A continuously differentiable function  $\alpha : [0, 1] \rightarrow \mathbb{R}$  is said to be a *lower solution* for problem (6) if the following properties hold:

- (i) there exists a unique function  $y_\alpha : [0, 1] \rightarrow \mathbb{R}$  such that

$$\begin{cases} y < y_\alpha(t) & \Rightarrow & f(t, y) < \alpha'(t), \\ y > y_\alpha(t) & \Rightarrow & f(t, y) > \alpha'(t); \end{cases} \tag{10}$$

- (ii)  $y_\alpha$  is continuously differentiable, and

$$(a(t)y_\alpha(t))' \geq g(t, \alpha(t)), \quad \text{for every } t \in [0, 1]; \tag{11}$$

- (iii)  $y_\alpha(0) \geq 0$  and  $\alpha(1) \sin \theta + y_\alpha(1) \cos \theta \leq 0$ .

**Definition 7.** A continuously differentiable function  $\beta : [0, 1] \rightarrow \mathbb{R}$  is said to be an *upper solution* for problem (6) if the following properties hold:




Figure 1. An illustration of the definition of lower and upper solutions from a dynamical point of view. Horizontal arrows represent the relative velocity  $x'$  of solutions of system (3) compared with  $\alpha'$  and  $\beta'$ , as stated in (10) and (12). Curved arrows indicate the essence of conditions (11) and (13), comparing  $y'$  with  $y'_\alpha$  and  $y'_\beta$

(j) there exists a unique function  $y_\beta : [0, 1] \rightarrow \mathbb{R}$  such that

$$\begin{cases} y < y_\beta(t) & \Rightarrow & f(t, y) < \beta'(t), \\ y > y_\beta(t) & \Rightarrow & f(t, y) > \beta'(t); \end{cases} \tag{12}$$

(jj)  $y_\beta$  is continuously differentiable, and

$$(a(t)y_\beta(t))' \leq g(t, \beta(t)), \quad \text{for every } t \in [0, 1]; \tag{13}$$

(jjj)  $y_\beta(0) \leq 0$  and  $\beta(1) \sin \theta + y_\beta(1) \cos \theta \geq 0$ .

For an intuitive meaning of the previous definitions, see Fig. 1.

### 2.2. Phase Plane Estimates

In this section, we provide some results which will be later used to prove the main theorems.

**Proposition 8.** *Suppose that  $a : [0, 1] \rightarrow \mathbb{R}$  satisfies (A1), (A2) and (A3). Then, there exists  $C > 0$  such that*

$$\int_0^t a(s)ds \leq Ca(t), \quad \text{for every } t \in [0, 1].$$

On the other hand, if  $a : [0, 1] \rightarrow \mathbb{R}$  satisfies (A1), (A2)' and (A3)', then there exists  $C > 0$  such that

$$\int_0^t a(s)ds \leq Ca(t), \quad \text{for every } t \in [0, \frac{1}{2}],$$

$$\int_t^1 a(s)ds \leq Ca(t), \quad \text{forevery } t \in [\frac{1}{2}, 1].$$

*Proof.* Assume  $a$  satisfies (A1), (A2)' and (A3)', the former case being easier. Consider the functions  $\psi_1 : [0, 1[ \rightarrow \mathbb{R}$  and  $\psi_2 : ]0, 1] \rightarrow \mathbb{R}$  defined by

$$\psi_1(t) = \begin{cases} \frac{\int_0^t a(s)ds}{a(t)} & t > 0, \\ 0 & t = 0; \end{cases} \quad \psi_2(t) = \begin{cases} \frac{\int_t^1 a(s)ds}{a(t)} & t < 1, \\ 0 & t = 1. \end{cases}$$

These functions are continuous, since by (A3)', we have

$$\psi_1(t) \leq t \quad \text{for all } t \in [0, \rho_0]; \quad \psi_2(t) \leq 1 - t \quad \text{for all } t \in [\rho_1, 1].$$

Then, the conclusion easily follows. □

We set

$$M = \max\{\|\hat{g}(t, x)\| : 0 \leq t \leq 1, \alpha(t) \leq x \leq \beta(t)\}, \tag{14}$$

where  $\hat{g}$  was defined in (7), and take a constant  $K$  satisfying

$$K > \max\{\|\alpha'\|_\infty, \|\beta'\|_\infty, \|y_\alpha\|_\infty, \|y_\beta\|_\infty, CM\},$$

where  $C$  is the constant introduced in Proposition 8.

We first modify the functions  $f(t, y)$  and  $g(t, x)$  as follows. Define  $\tilde{g} : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  by

$$\tilde{g}(t, x) = \begin{cases} g(t, \alpha(t)) + a(t)(x - \alpha(t)), & \text{if } x < \alpha(t), \\ g(t, x), & \text{if } \alpha(t) \leq x \leq \beta(t), \\ g(t, \beta(t)) + a(t)(x - \beta(t)), & \text{if } x > \beta(t), \end{cases}$$

and  $\tilde{f} : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  by

$$\tilde{f}(t, y) = \begin{cases} y, & \text{if } y \leq -K - 1, \\ f(t, y)(1 + K + y) - y(y + K), & \text{if } -K - 1 < y < -K, \\ f(t, y), & \text{if } -K \leq y \leq K, \\ f(t, y)(1 + K - y) + y(y - K), & \text{if } K < y < K + 1, \\ y, & \text{if } y \geq K + 1. \end{cases}$$

Let us consider the correspondingly modified problem

$$\begin{cases} x' = \tilde{f}(t, y), & (a(t)y)' = \tilde{g}(t, x), \\ y(0) = 0 = x(1) \sin \theta + y(1) \cos \theta. \end{cases} \tag{P̃}$$

We shall prove the existence of a solution of (P̃) and then verify that such a solution is indeed a solution of problem (6). To this aim, we define some

regions in the space  $[0, 1] \times \mathbb{R} \times \mathbb{R}$  and prove some invariance properties of them with respect to the solutions of the planar system

$$x' = \tilde{f}(t, y), \quad (a(t)y)' = \tilde{g}(t, x). \tag{\tilde{S}}$$

We set

$$\begin{aligned} A_{NE} &= \{(t, x, y) : t \in [0, 1], x > \beta(t), y > y_\beta(t)\}, \\ A_{SE} &= \{(t, x, y) : t \in [0, 1], x > \beta(t), y < y_\beta(t)\}, \\ A_{SW} &= \{(t, x, y) : t \in [0, 1], x < \alpha(t), y < y_\alpha(t)\}, \\ A_{NW} &= \{(t, x, y) : t \in [0, 1], x < \alpha(t), y > y_\alpha(t)\}. \end{aligned} \tag{15}$$

**Lemma 9.** *Let  $(x, y)$  be a solution of  $(\tilde{S})$  defined at a point  $t_0 \in [0, 1]$ . We have:*

- (i) *if  $y(t_0) > y_\beta(t_0)$ , then  $x'(t_0) > \beta'(t_0)$ ;*
- (ii) *if  $y(t_0) < y_\beta(t_0)$ , then  $x'(t_0) < \beta'(t_0)$ ;*
- (iii) *if  $y(t_0) > y_\alpha(t_0)$ , then  $x'(t_0) > \alpha'(t_0)$ ;*
- (iv) *if  $y(t_0) < y_\alpha(t_0)$ , then  $x'(t_0) < \alpha'(t_0)$ .*

*Proof.* We only prove (i), as the other assertions follow in a similar way. Assume  $y(t_0) > y_\beta(t_0)$ . Note that, as  $\|y_\beta\|_\infty \leq K$ , we have  $-K \leq y_\beta(t_0) < y(t_0)$ .

Suppose first that  $y(t_0) \leq K$ . Then,  $\tilde{f}(t_0, y(t_0)) = f(t_0, y(t_0))$  and hence, from (12), we get

$$(x - \beta)'(t_0) = f(t_0, y(t_0)) - \beta'(t_0) > 0.$$

Suppose next that  $K < y(t_0) < K + 1$ . Then, using (12) again and the fact that  $\|\beta'\|_\infty < K$ , we obtain

$$\begin{aligned} (x - \beta)'(t_0) &= f(t_0, y(t_0))(1 + K - y(t_0)) + y(t_0)(y(t_0) - K) - \beta'(t_0) \\ &> \beta'(t_0)(1 + K - y(t_0)) + K(y(t_0) - K) - \beta'(t_0) \\ &= \beta'(t_0)(K - y(t_0)) + K(y(t_0) - K) \\ &= (K - \beta'(t_0))(y(t_0) - K) > 0. \end{aligned}$$

Suppose finally that  $y(t_0) \geq K + 1$ . Then,

$$(x - \beta)'(t_0) = y(t_0) - \beta'(t_0) \geq K + 1 - \beta'(t_0) > 0.$$

Therefore,  $x'(t_0) > \beta'(t_0)$ . □

**Lemma 10.** *Let  $(x, y)$  be a solution of  $(\tilde{S})$  defined at a point  $t_0 \in [0, 1]$ , and suppose that both  $x(t_0) > \beta(t_0)$  and  $y(t_0) = y_\beta(t_0)$  hold. We have:*

- (i) *if  $t_0 \in ]0, 1[$ , then  $y'(t_0) > y'_\beta(t_0)$  ;*
- (ii) *if  $t_0 = 0$ , then there exists  $\delta > 0$  such that  $y(t) > y_\beta(t)$  for all  $t \in ]0, \delta[$ ;*
- (iii) *if  $t_0 = 1$ , then there exists  $\delta > 0$  such that  $y(t) < y_\beta(t)$  for all  $t \in ]1 - \delta, 1[$ .*

*Proof.* We first consider case (i). We recall that, from (13), we have

$$(a(t_0)y_\beta(t_0))' \leq g(t_0, \beta(t_0)).$$

Furthermore, we compute

$$\begin{aligned} (a(t)(y(t) - y_\beta(t)))' |_{t=t_0} &= a'(t_0)(y(t_0) - y_\beta(t_0)) + a(t_0)(y'(t_0) - y'_\beta(t_0)) \\ &= a(t_0)(y'(t_0) - y'_\beta(t_0)). \end{aligned} \tag{16}$$

Since  $x(t_0) > \beta(t_0)$ , we have  $\tilde{g}(t_0, x(t_0)) = g(t_0, \beta(t_0)) + a(t_0)(x(t_0) - \beta(t_0))$ , hence we obtain, using (16),

$$\begin{aligned} a(t_0)(y'(t_0) - y'_\beta(t_0)) &= (a(t_0)y(t_0))' - (a(t_0)y_\beta(t_0))' \\ &\geq \tilde{g}(t_0, x(t_0)) - g(t_0, \beta(t_0)) = a(t_0)(x(t_0) - \beta(t_0)). \end{aligned}$$

Since  $a(t_0) > 0$ , we conclude that  $y'(t_0) - y'_\beta(t_0) \geq x(t_0) - \beta(t_0) > 0$ .

We consider now case (ii). Let us set  $z(t) = a(t)(y(t) - y_\beta(t))$ . Since  $x(0) > \beta(0)$ , there exists  $\delta > 0$  such that  $x(s) > \beta(s)$ , for all  $s \in [0, \delta[$ . Pick  $t \in ]0, \delta[$ . Then, we have

$$\begin{aligned} z(t) &= \int_0^t z'(s)ds = \int_0^t (a(s)(y(s) - y_\beta(s)))' ds \\ &\geq \int_0^t (\tilde{g}(s, x(s)) - g(s, \beta(s))) ds = \int_0^t a(s)(x(s) - \beta(s)) ds > 0, \end{aligned}$$

hence  $y(t) - y_\beta(t) > 0$  for all  $t \in ]0, \delta[$ .

Case (iii) can be proved in a similar way. □

The following symmetric result can be proved similarly for the lower solution  $\alpha$ .

**Lemma 11.** *Let  $(x, y)$  be a solution of  $(\tilde{S})$  defined at a point  $t_0 \in [0, 1]$ , and suppose that both  $x(t_0) < \alpha(t_0)$  and  $y(t_0) = y_\alpha(t_0)$ . We have:*

- (i) *if  $t_0 \in ]0, 1[$ , then  $y'(t_0) < y'_\alpha(t_0)$ ;*
- (ii) *if  $t_0 = 0$ , then there exists  $\delta > 0$  such that  $y(t) < y_\alpha(t)$  for all  $t \in ]0, \delta[$ ;*
- (iii) *if  $t_0 = 1$ , then there exists  $\delta > 0$  such that  $y(t) > y_\alpha(t)$  for all  $t \in ]1 - \delta, 1[$ .*

The previous results allow us to prove some invariance properties of the regions  $A_{NE}, A_{SE}, A_{NW}, A_{SW}$  introduced in (15). To this aim, in the following statement, we consider a solution  $(x, y)$  of  $(\tilde{S})$  defined on a maximal interval of existence  $\mathcal{I}$ . Notice that, due to the linear growth of the functions  $\tilde{f}$  and  $\tilde{g}$ , if (A1), (A2) and (A3) hold, we can have the following two alternatives:

$$\mathcal{I} = [0, 1], \quad \mathcal{I} = ]0, 1].$$

On the other hand, if (A1), (A2)' and (A3)' hold, we can have the following four alternatives:

$$\mathcal{I} = [0, 1], \quad \mathcal{I} = ]0, 1], \quad \mathcal{I} = ]0, 1[, \quad \mathcal{I} = [0, 1[.$$

We use the conventional notation  $[s, s] = \{s\}$ , whenever  $s \in \mathbb{R}$ .

**Lemma 12.** *Let  $(x, y) : \mathcal{I} \rightarrow \mathbb{R}^2$  be a solution of  $(\tilde{S})$  defined at a point  $t_0 \in [0, 1]$ . We have:*

- (i) *if  $(t_0, x(t_0), y(t_0)) \in A_{NE}$ , then  $(t, x(t), y(t)) \in A_{NE}$  for all  $t \in [t_0, 1] \cap \mathcal{I}$ ;*



- (ii) if  $(t_0, x(t_0), y(t_0)) \in A_{SE}$ , then  $(t, x(t), y(t)) \in A_{SE}$  for all  $t \in [0, t_0] \cap \mathcal{I}$ ;
- (iii) if  $(t_0, x(t_0), y(t_0)) \in A_{SW}$ , then  $(t, x(t), y(t)) \in A_{SW}$  for all  $t \in [t_0, 1] \cap \mathcal{I}$ ;
- (iv) if  $(t_0, x(t_0), y(t_0)) \in A_{NW}$ , then  $(t, x(t), y(t)) \in A_{NW}$  for all  $t \in [0, t_0] \cap \mathcal{I}$ .

*Proof.* Let us prove the first assertion, the others follow similarly.

Let  $(t_0, x(t_0), y(t_0)) \in A_{NE}$  for some  $t_0 \in [0, 1]$ . By contradiction, assume that there exists  $t_1 \in ]t_0, 1] \cap \mathcal{I}$  such that  $(t, x(t), y(t)) \in A_{NE}$ , for every  $t \in [t_0, t_1[$ , and  $(t_1, x(t_1), y(t_1)) \notin A_{NE}$ . In particular we have either  $x(t_1) = \beta(t_1)$ , or  $y(t_1) = y_\beta(t_1)$ .

Since  $y(t_0) > y_\beta(t_0)$ , recalling Lemma 9, we have  $x'(t) > \beta'(t)$ , for every  $t \in [t_0, t_1[$ ; therefore, the first alternative is forbidden.

Finally, using Lemma 10 (i), we get a contradiction also in the case  $y(t_1) = y_\beta(t_1)$ . □

**Lemma 13.** *Let  $(x, y)$  be a solution of  $(\tilde{S})$ , defined on a nontrivial interval  $[t_1, t_2] \subseteq [0, 1]$ , satisfying  $\alpha(t_1) \leq x(t_1) \leq \beta(t_1)$  and  $\alpha(t_2) \leq x(t_2) \leq \beta(t_2)$ . Then,  $\alpha(t) \leq x(t) \leq \beta(t)$ , for all  $t \in [t_1, t_2]$ .*

*Proof.* We assume, by contradiction, that there exists  $t_0 \in ]t_1, t_2[$  with  $x(t_0) > \beta(t_0)$ .

Suppose first that  $y(t_0) > y_\beta(t_0)$ . Then  $(t_0, x(t_0), y(t_0)) \in A_{NE}$ . From Lemma 12, we have  $(t_2, x(t_2), y(t_2)) \in A_{NE}$ . In particular,  $x(t_2) > \beta(t_2)$ , a contradiction.

Suppose now that  $y(t_0) = y_\beta(t_0)$ . From Lemma 10, we see that  $y(t) > y_\beta(t)$  in a right neighbourhood of  $t_0$ , and we conclude as before.

Finally, if  $y(t_0) < y_\beta(t_0)$ , we have  $(t_0, x(t_0), y(t_0)) \in A_{SE}$ . From Lemma 12, we have  $(t_1, x(t_1), y(t_1)) \in A_{SE}$ . In particular,  $x(t_1) > \beta(t_1)$ , again a contradiction.

Hence, we conclude that  $x(t) \leq \beta(t)$  for every  $t \in [t_1, t_2]$ .

In a similar way, we can prove that  $x(t) \geq \alpha(t)$  for every  $t \in [t_1, t_2]$ , thus concluding the proof. □

**Lemma 14.** *Let  $(x, y)$  be a solution of  $(\tilde{S})$ , defined on a nontrivial interval  $[0, t_2] \subseteq [0, 1]$ , satisfying the following properties:*

$$y(0) = 0, \quad \alpha(0) \leq x(0) \leq \beta(0), \quad \alpha(t_2) \leq x(t_2) \leq \beta(t_2).$$

*Then  $|y(t)| \leq K$ , for all  $t \in [0, t_2]$ .*

*Proof.* By Lemma 13, we have that  $\alpha(t) \leq x(t) \leq \beta(t)$ , for every  $t \in [0, t_2]$ . In particular,

$$\tilde{g}(t, x(t)) = g(t, x(t)) = a(t)\hat{g}(t, x(t)),$$

for all  $t \in [0, t_2]$ . Let us set  $z(t) = a(t)y(t)$ . Then, for all  $t \in [0, t_2]$ ,

$$z(t) = \int_0^t z'(s)ds = \int_0^t \tilde{g}(s, x(s))ds = \int_0^t a(s)\hat{g}(s, x(s))ds.$$

Recalling the definition (14) of  $M$  and Proposition 8, we deduce that

$$|z(t)| \leq \int_0^t Ma(s)ds \leq MCa(t).$$

Therefore,  $a(t)|y(t)| \leq MCa(t)$  for all  $t \in [0, t_2]$ . If  $a(t) \neq 0$ , we obtain

$$|y(t)| \leq MC < K,$$

for every  $t \in ]0, t_2]$ . This inequality is trivially satisfied also in case  $t = 0$ , hence the lemma is completely proved.  $\square$

Arguing similarly, we can prove the following result.

**Lemma 15.** *Assume (A1), (A2)' and (A3)'. Let  $(x, y)$  be a solution of  $(\tilde{S})$ , defined on a nontrivial interval  $[t_1, 1] \subseteq [0, 1]$ , satisfying the following properties:*

$$y(1) = 0, \quad \alpha(t_1) \leq x(t_1) \leq \beta(t_1), \quad \alpha(1) \leq x(1) \leq \beta(1).$$

*Then,  $|y(t)| \leq K$ , for all  $t \in [t_1, 1]$ .*

So far, we have proved the following a priori bounds.

**Proposition 16.** *Assume (A1), (A2) and (A3). If  $(x, y)$  is a solution of  $(\tilde{P})$ , satisfying  $\alpha(0) \leq x(0) \leq \beta(0)$  and  $\alpha(1) \leq x(1) \leq \beta(1)$ , then  $(x, y)$  is a solution of problem (6) and satisfies  $\alpha(t) \leq x(t) \leq \beta(t)$ , for all  $t \in [0, 1]$ .*

*Proof.* It is an immediate consequence of the application of Lemma 13 with  $[t_1, t_2] = [0, 1]$  and Lemma 14 with  $[0, t_2] = [0, 1]$ .  $\square$

**Proposition 17.** *Assume (A1), (A2)' and (A3)'. If  $(x, y)$  is a solution of  $(\tilde{P})$ , with  $\theta = 0$ , satisfying, for a certain  $t_0 \in ]0, 1[$ ,*

$$\alpha(0) \leq x(0) \leq \beta(0), \quad \alpha(t_0) \leq x(t_0) \leq \beta(t_0), \quad \alpha(1) \leq x(1) \leq \beta(1);$$

*then  $(x, y)$  is a solution of problem (6), with  $\theta = 0$ , and satisfies  $\alpha(t) \leq x(t) \leq \beta(t)$ , for all  $t \in [0, 1]$ .*

*Proof.* We need to apply twice Lemma 13 with  $[t_1, t_2] = [0, t_0]$  and  $[t_1, t_2] = [t_0, 1]$ . Then we apply Lemma 14 with  $[0, t_2] = [0, t_0]$  and Lemma 15 with  $[t_1, 1] = [t_0, 1]$ .  $\square$

Summing up, to prove Theorem 2, we need to find a solution of  $(\tilde{P})$  satisfying the assumptions of Proposition 16. Similarly, to prove Theorem 4, we need to find a solution of  $(\tilde{P})$ , with  $\theta = 0$ , satisfying the assumptions of Proposition 17.

### 3. Proof of the Theorems

#### 3.1. Proof of Theorem 2

To prove our result, we shall apply a shooting argument, with the aim of finding  $\sigma \in \mathbb{R}$  such that the solution  $(x, y)$  of the Cauchy problem

$$\begin{cases} x' = \tilde{f}(t, y), & (a(t)y)' = \tilde{g}(t, x), \\ x(0) = \sigma, & y(0) = 0 \end{cases} \tag{17}$$

also satisfies  $x(1) \sin \theta + y(1) \cos \theta = 0$ .

We start by defining the flow associated with system  $(\tilde{S})$ . Let  $\mathcal{X}$  be the set of initial data

$$\mathcal{X} = \{(t_0, \sigma, \tau) \in [0, 1] \times \mathbb{R}^2 : \tau = 0 \text{ if } t_0 = 0\},$$

and consider the solution

$$(x(\cdot), y(\cdot)) = \Phi(\cdot; t_0, \sigma, \tau) = (\Phi_1(\cdot; t_0, \sigma, \tau), \Phi_2(\cdot; t_0, \sigma, \tau))$$

of  $(\tilde{S})$  satisfying  $x(t_0) = \sigma$  and  $y(t_0) = \tau$ . The proof concerning the existence of such a solution, which is not completely standard due to the presence of the singularity at 0, is given in the Appendix. This solution will be proved to be defined on  $]0, 1]$ , thanks to the linear growth of the functions  $\tilde{f}$  and  $\tilde{g}$ , but not necessarily at  $t = 0$ . However, if  $t_0 = 0$ , the solution is defined on the whole interval  $[0, 1]$ . We denote by  $\mathcal{D} \subseteq \mathbb{R}^4$  the domain of the flow  $\Phi = \Phi(t; t_0, \sigma, \tau)$ . We have

$$]0, 1] \times \mathcal{X} \subseteq \mathcal{D} \subseteq [0, 1] \times \mathcal{X}.$$

The continuity of the flow  $\Phi$  follows from the continuous dependence of the solutions of  $(\tilde{S})$  on the initial data. Again, the proof of this fact is given in the Appendix.

Let us fix  $U_0 > 0$  such that

$$-U_0 < \alpha(0) \leq \beta(0) < U_0, \tag{18}$$

and define the continuous curve

$$\begin{aligned} \mathcal{C} &: [-U_0, U_0] \rightarrow \mathbb{R}^2 \\ \mathcal{C}(\sigma) &= (x_{\mathcal{C}}(\sigma), y_{\mathcal{C}}(\sigma)) := \Phi(1; 0, \sigma, 0). \end{aligned} \tag{19}$$

The following proposition localises the curve  $\mathcal{C}$ .

**Proposition 18.** *Let  $\mathcal{C}$  be the curve defined by (19). Then, the following properties hold:*

- (i)  $(1, \mathcal{C}(\sigma)) \in A_{SW}$  for every  $\sigma \in [-U_0, \alpha(0)[$ ;
- (ii)  $(1, \mathcal{C}(\sigma)) \in A_{NE}$  for every  $\sigma \in ]\beta(0), U_0]$ ;
- (iii)  $(1, \mathcal{C}(\sigma)) \notin A_{NW} \cup A_{SE}$  for all  $\sigma \in [-U_0, U_0]$ .

*Proof.* Let us prove (i). Let  $(x, y)$  be the solution of  $(\tilde{S})$  satisfying  $x(0) = \sigma < \alpha(0)$  and  $y(0) = 0$ . Recall that, by the definition of lower solution,  $y_{\alpha}(0) \geq 0$ . Assume first that  $y_{\alpha}(0) > 0$ . Then  $(0, x(0), y(0)) \in A_{SW}$ . By Lemma 12, we have that  $(t, x(t), y(t)) \in A_{SW}$  for all  $t \in [0, 1]$  and, in particular,  $(1, \mathcal{C}(\sigma)) = (1, x(1), y(1)) \in A_{SW}$ .

Assume next that  $y_{\alpha}(0) = 0$ . Then, by Lemma 11, there exists  $\delta > 0$  such that  $y(t) < y_{\alpha}(t)$  for all  $t \in ]0, \delta[$ . By continuity, we can find  $t_1 \in ]0, \delta[$  such that  $(t_1, x(t_1), y(t_1)) \in A_{SW}$ . Therefore, by Lemma 12, we have  $(t, x(t), y(t)) \in A_{SW}$  for all  $t \in [t_1, 1]$  and, in particular,  $(1, \mathcal{C}(\sigma)) \in A_{SW}$ .

The proof of (ii) is similar, hence we omit it, for brevity.

Let us prove (iii). Suppose, by contradiction, that there is  $\sigma \in [-U_0, U_0]$  such that  $(1, \mathcal{C}(\sigma)) \in A_{NW} \cup A_{SE}$ . Let  $(x, y)$  be the solution of  $(\tilde{S})$  satisfying  $x(0) = \sigma$  and  $y(0) = 0$ .

Suppose first that  $(1, \mathcal{C}(\sigma)) \in A_{NW}$ . Since  $(1, x(1), y(1)) \in A_{NW}$ , by Lemma 12 we have that  $(t, x(t), y(t)) \in A_{NW}$  for all  $t \in [0, 1]$  and, in particular,  $(0, \sigma, 0) \in A_{NW}$ . This is a contradiction, since any point  $(0, \sigma, y) \in A_{NW}$  satisfies  $y > y_\alpha(0) \geq 0$ .

Suppose now that  $(1, \mathcal{C}(\sigma)) \in A_{SE}$ . Then, by Lemma 12 again, we have that  $(t, x(t), y(t)) \in A_{SE}$  for all  $t \in [0, 1]$  and, in particular,  $(0, \sigma, 0) \in A_{SE}$ . This is a contradiction, since any  $(0, \sigma, y) \in A_{SE}$  satisfies  $y < y_\beta(0) \leq 0$ .  $\square$

We shall consider the restriction of  $\mathcal{C}$  on some intervals  $[\sigma_\ell, \sigma_r]$  so that  $\alpha(1) \leq x_\mathcal{C}(\sigma) \leq \beta(1)$  for all  $\sigma \in [\sigma_\ell, \sigma_r]$ . To this aim, we set

$$\begin{aligned} \sigma_\ell &= \min\{\sigma \in [-U_0, U_0] : x_\mathcal{C}(s) \geq \alpha(1) \text{ for all } s \in [\sigma, U_0]\}; \\ \sigma_r &= \max\{\sigma \in [-U_0, U_0] : x_\mathcal{C}(s) \leq \beta(1) \text{ for all } s \in [-U_0, \sigma]\}. \end{aligned}$$

Observe that, from Proposition 18 (i)–(ii), we have

$$\alpha(0) \leq \sigma_\ell \leq \sigma_r \leq \beta(0), \tag{20}$$

and

$$x_\mathcal{C}(\sigma_\ell) = \alpha(1), \quad x_\mathcal{C}(\sigma_r) = \beta(1).$$

Then, by Proposition 18 (iii), we have

$$y_\mathcal{C}(\sigma_\ell) \leq y_\alpha(1), \quad y_\mathcal{C}(\sigma_r) \geq y_\beta(1).$$

Since  $\cos \theta \geq 0$ , we get both

$$x_\mathcal{C}(\sigma_\ell) \sin \theta + y_\mathcal{C}(\sigma_\ell) \cos \theta \leq \alpha(1) \sin \theta + y_\alpha(1) \cos \theta \leq 0,$$

and

$$x_\mathcal{C}(\sigma_r) \sin \theta + y_\mathcal{C}(\sigma_r) \cos \theta \geq \beta(1) \sin \theta + y_\beta(1) \cos \theta \geq 0.$$

Since the curve  $\mathcal{C}$  is continuous, we can find  $\sigma \in [\sigma_\ell, \sigma_r]$  such that

$$x_\mathcal{C}(\sigma) \sin \theta + y_\mathcal{C}(\sigma) \cos \theta = 0.$$

Therefore, the function  $(x, y) = \Phi(\cdot; 0, \sigma, 0)$  is a solution of problem  $(\tilde{P})$ . Notice that we have both  $\alpha(0) \leq x(0) = \sigma \leq \beta(0)$ , from (20), and  $\alpha(1) \leq x(1) = x_\mathcal{C}(\sigma) \leq \beta(1)$ , from the definition of the interval  $[\sigma_\ell, \sigma_r]$ . By Proposition 16,  $(x, y)$  is a solution of problem (6) and satisfies  $\alpha(t) \leq x(t) \leq \beta(t)$ , for all  $t \in [0, 1]$ . The proof of Theorem 2 is, thus, completed.

### 3.2. Proof of Theorem 4

To prove our second result, we shall apply a *double shooting* argument, with the aim of finding  $\sigma \in \mathbb{R}$  such that the solution  $(x, y)$  of the Cauchy problem (17) also satisfies  $y(1) = 0$ .

To define the flow associated with system  $(\tilde{S})$  under assumptions (A1), (A2)' and (A3)', the set of possible initial data is now

$$\mathcal{X} = \{(t_0, \sigma, \tau) \in [0, 1] \times \mathbb{R}^2 : \tau = 0 \text{ if } t_0 = 0 \text{ or } t_0 = 1\}.$$

The solutions are defined on  $]0, 1[$  but not necessarily at  $t = 0$  or  $t = 1$ . See the Appendix for details. We denote by  $\mathcal{D} \subseteq \mathbb{R}^4$  the domain of the flow  $\Phi = \Phi(t; t_0, \sigma, \tau)$ . We have

$$]0, 1[ \times \mathcal{X} \subseteq \mathcal{D} \subseteq [0, 1] \times \mathcal{X}.$$

Let us fix  $U_0 > 0$  such that

$$-U_0 < \min\{\alpha(0), \alpha(1)\} \leq \max\{\beta(0), \beta(1)\} < U_0.$$

For any  $\sigma_0, \sigma_1 \in [-U_0, U_0]$ , we consider the the initial value problem

$$\begin{cases} x' = \tilde{f}(t, y), & (a(t)y)' = \tilde{g}(t, x), \\ x(0) = \sigma_0, & y(0) = 0, \end{cases} \tag{21}$$

and the final value problem

$$\begin{cases} x' = \tilde{f}(t, y), & (a(t)y)' = \tilde{g}(t, x), \\ x(1) = \sigma_1, & y(1) = 0. \end{cases} \tag{22}$$

We use a shooting argument to find a solution  $(x_{\sigma_0}, y_{\sigma_0})$  of (21) (defined on  $[0, 1[$ ), and a solution  $(x^{\sigma_1}, y^{\sigma_1})$  of (22) (defined on  $]0, 1]$ ), satisfying

$$(x_{\sigma_0}(\frac{1}{2}), y_{\sigma_0}(\frac{1}{2})) = (x^{\sigma_1}(\frac{1}{2}), y^{\sigma_1}(\frac{1}{2})).$$

Clearly, the function  $(x, y)$  is defined by

$$(x(t), y(t)) = \begin{cases} (x_{\sigma_0}(t), y_{\sigma_0}(t)), & \text{if } 0 \leq t \leq \frac{1}{2}, \\ (x^{\sigma_1}(t), y^{\sigma_1}(t)), & \text{if } \frac{1}{2} < t \leq 1, \end{cases} \tag{23}$$

will be the solution of  $(\tilde{P})$  we are looking for.

Let us define two continuous curves  $\mathcal{C}_0, \mathcal{C}_1 : [-U_0, U_0] \rightarrow \mathbb{R}^2$  by

$$\begin{aligned} \mathcal{C}_0(\sigma) &= (x_{\mathcal{C}}^0(\sigma), y_{\mathcal{C}}^0(\sigma)) := \Phi(\frac{1}{2}; 0, \sigma, 0), \\ \mathcal{C}_1(\sigma) &= (x_{\mathcal{C}}^1(\sigma), y_{\mathcal{C}}^1(\sigma)) := \Phi(\frac{1}{2}; 1, \sigma, 0). \end{aligned} \tag{24}$$

The following statement describes some localisation properties of the curves  $\mathcal{C}_0$  and  $\mathcal{C}_1$ .

**Proposition 19.** *Let  $\mathcal{C}_0$  and  $\mathcal{C}_1$  be the curves defined by (24). Then, the following properties hold:*

- (i)  $(\frac{1}{2}, \mathcal{C}_0(\sigma)) \in A_{SW}$  and  $(\frac{1}{2}, \mathcal{C}_1(\sigma)) \in A_{NW}$  for every  $\sigma \in [-U_0, \alpha(0)[$ ;
- (ii)  $(\frac{1}{2}, \mathcal{C}_0(\sigma)) \in A_{NE}$  and  $(\frac{1}{2}, \mathcal{C}_1(\sigma)) \in A_{SE}$  for every  $\sigma \in ]\beta(0), U_0]$ ;
- (iii)  $(\frac{1}{2}, \mathcal{C}_0(\sigma)) \notin A_{NW} \cup A_{SE}$  for all  $\sigma \in [-U_0, U_0]$ ;
- (iv)  $(\frac{1}{2}, \mathcal{C}_1(\sigma)) \notin A_{SW} \cup A_{NE}$  for all  $\sigma \in [-U_0, U_0]$ .

The proof can be adapted from that of Propositions 18. We prove now that the two curves have a common value.

**Proposition 20.** *Let  $\mathcal{C}_0$  and  $\mathcal{C}_1$  be the curves defined by (24). Then, there are  $\sigma_0, \sigma_1 \in ]-U_0, U_0[$  such that  $\mathcal{C}_0(\sigma_0) = \mathcal{C}_1(\sigma_1)$ .*

*Proof.* We shall consider the restriction of  $\mathcal{C}_0$  and  $\mathcal{C}_1$  on some intervals  $[\sigma_\ell^0, \sigma_r^0]$  and  $[\sigma_\ell^1, \sigma_r^1]$ , respectively, so that  $\alpha(\frac{1}{2}) \leq x_{\mathcal{C}}^0(\sigma) \leq \beta(\frac{1}{2})$  for all  $\sigma \in [\sigma_\ell^0, \sigma_r^0]$  and  $\alpha(\frac{1}{2}) \leq x_{\mathcal{C}}^1(\sigma) \leq \beta(\frac{1}{2})$  for all  $\sigma \in [\sigma_\ell^1, \sigma_r^1]$ . To this aim, we set

$$\begin{aligned} \sigma_\ell^0 &= \min\{\sigma \in [-U_0, U_0] : x_{\mathcal{C}}^0(s) \geq \alpha(\frac{1}{2}) \text{ for all } s \in [\sigma, U_0]\}; \\ \sigma_r^0 &= \max\{\sigma \in [-U_0, U_0] : x_{\mathcal{C}}^0(s) \leq \beta(\frac{1}{2}) \text{ for all } s \in [-U_0, \sigma]\}; \\ \sigma_\ell^1 &= \min\{\sigma \in [-U_0, U_0] : x_{\mathcal{C}}^1(s) \geq \alpha(\frac{1}{2}) \text{ for all } s \in [\sigma, U_0]\}; \\ \sigma_r^1 &= \max\{\sigma \in [-U_0, U_0] : x_{\mathcal{C}}^1(s) \leq \beta(\frac{1}{2}) \text{ for all } s \in [-U_0, \sigma]\}. \end{aligned}$$

Observe that, from Proposition 19 (i)–(ii), we have

$$\alpha(0) \leq \sigma_\ell^0 \leq \sigma_r^0 \leq \beta(0) \quad \text{and} \quad \alpha(1) \leq \sigma_\ell^1 \leq \sigma_r^1 \leq \beta(1). \tag{25}$$

Moreover,

$$x_\ell^0(\sigma_\ell^0) = \alpha(\frac{1}{2}) = x_\ell^1(\sigma_\ell^1) \quad \text{and} \quad x_r^0(\sigma_r^0) = \beta(\frac{1}{2}) = x_r^1(\sigma_r^1).$$

By Proposition 19 (iii)–(iv), we have

$$y_\ell^0(\sigma_\ell^0) \leq y_\alpha(\frac{1}{2}) \leq y_\ell^1(\sigma_\ell^1) \quad \text{and} \quad y_r^0(\sigma_r^0) \geq y_\beta(\frac{1}{2}) \geq y_r^1(\sigma_r^1).$$

Since the curves are continuous, they must cross each other at some point  $(x_\ell^0(\sigma_0), y_\ell^0(\sigma_0)) = (x_\ell^1(\sigma_1), y_\ell^1(\sigma_1))$ , with  $\sigma_0 \in [\sigma_\ell^0, \sigma_r^0]$  and  $\sigma_1 \in [\sigma_\ell^1, \sigma_r^1]$ .  $\square$

The parameters  $\sigma_0, \sigma_1$  obtained in Proposition 20 permit us to define the solution  $(x, y)$  of problem  $(\tilde{P})$  as in (23). In particular, we have

$$\alpha(0) \leq x(0) = \sigma_0 \leq \beta(0), \quad \alpha(1) \leq x(1) = \sigma_1 \leq \beta(1),$$

from (25), and

$$\alpha(\frac{1}{2}) \leq x(\frac{1}{2}) \leq \beta(\frac{1}{2}),$$

from the definition of the intervals  $[\sigma_\ell^0, \sigma_r^0]$  and  $[\sigma_\ell^1, \sigma_r^1]$ . By Proposition 17,  $(x, y)$  is a solution of problem (6), with  $\theta = 0$ , and satisfies  $\alpha(t) \leq x(t) \leq \beta(t)$ , for all  $t \in [0, 1]$ . The proof of Theorem 4 is, thus, completed.

### 4. Examples and Final Remarks

In this section, we provide some possible applications of our theorems.

In (4), we have considered for simplicity a differential equation ruled by a weighted  $p$ -Laplacian. In a similar way, we can consider a *double-weighted*  $\phi$ -Laplace equation, in the unitary ball, of the type

$$\operatorname{div}\left(\eta(|x|)\phi(m(|x|)\nabla v(x))\right) = h(|x|, v(x)), \tag{26}$$

where  $\eta, m : [0, 1] \rightarrow \mathbb{R}^+$  are positive continuous functions,  $\phi(w) = \psi(|w|)\frac{w}{|w|}$ , being  $\psi : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  an odd increasing diffeomorphism, and  $h : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  is continuous and locally Lipschitz continuous with respect to the second variable. In the case of Eq. (4), we have  $m \equiv 1$  and  $\psi(y) = |y|^{p-2}$ . Another example is the relativistic curvature operator provided by

$$\psi(y) = \frac{y}{\sqrt{1 - y^2}},$$

cf. [1], where  $I = ] - 1, 1[$ . The study of radial solutions of Eq. (26) leads to the equivalent equation

$$(r^{N-1}\eta(r)\psi(m(r)u'))' = r^{N-1}h(r, u), \quad r \in [0, 1],$$

which can be written as a planar system of the form

$$x' = \omega(t)\psi^{-1}(y), \quad (t^{N-1}\eta(t)y)' = t^{N-1}h(t, x),$$

where  $\omega(t) = 1/m(t)$ , which is a special case of (3).

Theorem 2 may be applied to study the boundary value problem

$$\begin{cases} x' = \omega(t)\psi^{-1}(y), & (t^{N-1}\eta(t)y)' = t^{N-1}h(t, x), \\ y(0) = 0 = x(1) \sin \theta + y(1) \cos \theta. \end{cases} \tag{27}$$

Notice that all the regularity assumptions required in Theorem 2 immediately hold. So, if we are able to provide a well-ordered couple of lower/upper solutions for problem (27), then we can successfully apply it.

The following statement describes a possible example of application in the case of *constant* lower and upper solutions.

**Corollary 21.** *Let  $\theta \in [0, \frac{\pi}{2}]$ , and assume the existence of some constants  $\alpha \leq 0 \leq \beta$  such that  $h(t, \alpha) \leq 0 \leq h(t, \beta)$  for every  $t \in [0, 1]$ . Then, problem (27) has a solution  $(x, y)$  such that  $\alpha \leq x(t) \leq \beta$ , for every  $t \in [0, 1]$ .*

*Proof.* It is easy to verify that the constant functions  $\alpha$  and  $\beta$  fulfill the conditions in Definitions 6 and 7 with the choice  $y_\alpha = y_\beta \equiv 0$ . Then, Theorem 2 applies, thus completing the proof. □

In particular, the previous corollary permits us to find an existence result for Eq. (26) with Dirichlet or Neumann boundary conditions on the unitary ball  $\mathcal{B}$ .

**Corollary 22.** *Assume the existence of some constants  $\alpha \leq 0 \leq \beta$  such that  $h(r, \alpha) \leq 0 \leq h(r, \beta)$  for every  $r \in [0, 1]$ . Then, problem*

$$\begin{cases} \operatorname{div}\left(\eta(|x|)\phi(m(|x|)\nabla v(x))\right) = h(|x|, v(x)) & \text{in } \mathcal{B}, \\ v = 0 & \text{on } \partial\mathcal{B} \end{cases}$$

*has a solution  $v$  such that  $\alpha \leq v(x) \leq \beta$  for every  $x \in \overline{\mathcal{B}}$ .*

**Corollary 23.** *Assume the existence of some constants  $\alpha \leq 0 \leq \beta$  such that  $h(r, \alpha) \leq 0 \leq h(r, \beta)$  for every  $r \in [0, 1]$ . Then, problem*

$$\begin{cases} \operatorname{div}\left(\eta(|x|)\phi(m(|x|)\nabla v(x))\right) = h(|x|, v(x)) & \text{in } \mathcal{B}, \\ \partial_\nu v = 0 & \text{on } \partial\mathcal{B} \end{cases}$$

*has a solution  $v$  such that  $\alpha \leq v(x) \leq \beta$  for every  $x \in \overline{\mathcal{B}}$ .*

*Example 24.* Let us consider two locally Lipschitz continuous functions  $f, g : \mathbb{R} \rightarrow \mathbb{R}$  satisfying

$$yf(y) > 0, \quad \text{when } y \neq 0,$$

and

$$\liminf_{x \rightarrow -\infty} g(x) < -\bar{g}, \quad \limsup_{x \rightarrow +\infty} g(x) > \bar{g},$$

for some  $\bar{g} > 0$ . Then, given  $a(t)$  satisfying (A1), (A2) and (A3), for any function  $e : [0, 1] \rightarrow \mathbb{R}$  such that  $\|e\|_\infty \leq \bar{g}$ , the problem

$$\begin{cases} x' = f(y), & (a(t)y(t))' = a(t)[g(x) + e(t)], \\ y(0) = 0 = x(1) \sin \theta + y(1) \cos \theta, \end{cases}$$

with  $\theta \in [0, \frac{\pi}{2}]$ , has a solution by applying Corollary 21. As particular cases, choosing

$$p \leq 2 \leq q, \quad \frac{1}{p} + \frac{1}{q} = 1, \quad f(y) = |y|^{q-2}y, \quad a(t) = t^{N-1},$$

with  $N \geq 1$ , the problems in the unitary ball of  $\mathbb{R}^N$

$$\begin{cases} \Delta_p(v) = g(v) + e(|x|) & \text{in } \mathcal{B}, \\ v = 0 & \text{on } \partial\mathcal{B}, \end{cases} \quad \begin{cases} \Delta_p(v) = g(v) + e(|x|) & \text{in } \mathcal{B}, \\ \partial_\nu v = 0 & \text{on } \partial\mathcal{B}, \end{cases}$$

have at least one radial solution. We, thus, recover some well-known classical results.

Analogous considerations allow to generalize problem (9), providing further applications of Theorem 4.

*Example 25.* Let  $g(x)$  and  $e(t)$  be as in the previous example. We can apply Theorem 4 so to find a solution of the problem

$$\begin{cases} (\sin^{N-2}(\pi t)x')' = \sin^{N-2}(\pi t)[g(x) + e(t)], \\ x'(0) = 0 = x'(1). \end{cases}$$

*Remark 26.* In this paper, we have treated the case when the pair of lower and upper solutions is well ordered. The non-well-ordered case is left as an open problem.

**Author contributions** All authors contributed at the same level to the results in the paper.

**Funding** Open access funding provided by Università degli Studi di Trieste within the CRUI-CARE Agreement. International Science Program (ISP), Uppsala University.

**Data Availability Statement** Not applicable.

**Declarations**

**Conflict of interest** There are no competing interests.

**Ethical approval** Not applicable.

**Open Access.** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by



statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Appendix

In this appendix, we prove the continuity of the flow  $\Phi : \mathcal{D} \subseteq [0, 1] \times \mathcal{X} \rightarrow \mathbb{R}^2$ , introduced in Sect. 3.1, associated with system  $(\tilde{S})$ . Then, we will provide the corresponding result for the situation treated in Sect. 3.2.

Because of the presence of the singularity at  $t = 0$ , we provide a proof of existence, uniqueness and continuous dependence on initial data properties of the solutions of the Cauchy problems (17). Similar properties have been studied for second-order differential equations presenting a singularity. See e.g. [9, Appendix] or [2–4, 12].

We recall that, under the assumptions of Theorem 2, the function  $a : [0, 1] \rightarrow \mathbb{R}$  is of class  $C^1$ , positive in the interval  $]0, 1]$ , increasing in  $[0, \rho_0] \subseteq [0, 1]$ , and satisfies  $a(0) = 0$ . To simplify the notation, we consider the Cauchy problem

$$\begin{cases} u' = F(t, y), & (a(t)y)' = a(t)G(t, x), \\ u(0) = u_0, & y(0) = 0, \end{cases} \tag{28}$$

where  $\tilde{f}(t, y) = F(t, y)$  and  $\tilde{g}(t, x) = a(t)G(t, x)$ . Since, by construction, both  $F$  and  $G$  are locally Lipschitz continuous with respect to the second variable and they have an at most linear growth, we can assume that there exists  $A > 0$  such that

$$|F(t, y)| \leq A(1 + |y|), \quad |G(t, x)| \leq A(1 + |x|), \tag{29}$$

for every  $t \in [0, 1]$  and  $x, y \in \mathbb{R}$ .

At first we notice that, for  $t \in ]0, 1]$ , the differential system in (28) can be rewritten as

$$x' = F(t, y), \quad y' = -\frac{a'(t)}{a(t)}y + G(t, x),$$

thus obtaining a planar system for which we can easily verify local existence and uniqueness of the solutions for the Cauchy problems. Moreover, since (29) holds, such solutions are globally defined on  $]0, 1]$ .

Hence, in what follows, we focus our attention on Cauchy problems of the form

$$\begin{cases} x' = F(t, y), & (a(t)y)' = a(t)G(t, x), \\ x(0) = x_0, & y(0) = 0, \end{cases} \tag{30}$$

where  $x_0 \in \mathbb{R}$ . In particular, it will be sufficient to prove existence, uniqueness and continuous dependence on initial data for such Cauchy problems only in a right neighborhood of 0. Since the functions  $F$  and  $G$  satisfy (29), we can then easily recover these properties in the whole interval  $[0, 1]$ .

We start by stating the local existence and uniqueness theorem.

**Theorem 27.** *For every  $x_0 \in \mathbb{R}$  there exists  $\tau > 0$  such that there is a unique solution  $(x, y) : [0, \tau] \rightarrow \mathbb{R}^2$  of the Cauchy problem (30).*

*Proof.* Since the functions  $F$  and  $G$  are continuous and locally Lipschitz continuous with respect to the variables  $x, y$ , we can find constants  $\mathcal{M}$  and  $L$  such that, for every  $s \in [0, 1]$ ,

$$\begin{aligned} |y| \leq 1 &\Rightarrow |F(s, y)| \leq \mathcal{M}, \\ |x - x_0| \leq 1 &\Rightarrow |G(s, x)| \leq \mathcal{M}, \\ |y_1| \leq 1 \text{ and } |y_2| \leq 1 &\Rightarrow |F(s, y_1) - F(s, y_2)| \leq L|y_1 - y_2|, \\ |x_1 - x_0| \leq 1 \text{ and } |x_2 - x_0| \leq 1 &\Rightarrow |G(s, x_1) - G(s, x_2)| \leq L|x_1 - x_2|. \end{aligned}$$

Pick a constant  $\tau$  satisfying

$$\tau < \max \left\{ \rho_0, \frac{1}{\mathcal{M}}, \frac{1}{L} \right\},$$

and introduce the Banach space  $X = C^0([0, \tau], \mathbb{R}^2)$ , endowed with the norm  $\|(x, y)\|_X = \max\{\|x\|_\infty, \|y\|_\infty\}$ . Set

$$\mathcal{B} = \{(x, y) \in C^0([0, \tau], \mathbb{R}^2) : \|x - x_0\|_\infty \leq 1 \text{ and } \|y\|_\infty \leq 1\},$$

and define the function  $T : \mathcal{B} \rightarrow \mathcal{B}$  by

$$T(x, y)[t] = (T_1(x, y)[t], T_2(x, y)[t]),$$

where

$$T_1(x, y)[t] = x_0 + \int_0^t F(s, y(s))ds, \quad T_2(x, y)[t] = \frac{1}{a(t)} \int_0^t a(s)G(s, x(s))ds.$$

Notice that  $T(x, y) \in \mathcal{B}$ , for every  $(x, y) \in \mathcal{B}$ . Indeed, for every  $t \in ]0, \tau]$ , we have both

$$|T_1(x, y)[t] - x_0| = \left| \int_0^t F(s, y(s))ds \right| \leq t\mathcal{M} < 1,$$

and, recalling (A3),

$$\begin{aligned} |T_2(x, y)[t]| &= \left| \frac{1}{a(t)} \int_0^t a(s)G(s, x(s))ds \right| \\ &\leq \frac{1}{a(t)} \int_0^t a(s)\mathcal{M}ds \leq t\mathcal{M} < 1. \end{aligned}$$

Let us prove that the function  $T$  is a contraction. We set  $\kappa = \tau L < 1$ . Then, given any  $(x_1, y_1), (x_2, y_2) \in \mathcal{B}$  and for every  $t \in [0, \tau]$ , we have both

$$\begin{aligned} |T_1(x_1, y_1)[t] - T_1(x_2, y_2)[t]| &= \left| \int_0^t F(s, y_1(s)) - F(s, y_2(s))ds \right| \\ &\leq tL\|y_1 - y_2\|_\infty < \kappa\|(x_1, y_1) - (x_2, y_2)\|_X, \end{aligned}$$

and, recalling (A3),

$$\begin{aligned} |T_2(x_1, y_1)[t] - T_2(x_2, y_2)[t]| &= \left| \frac{1}{a(t)} \int_0^t a(s)(G(s, x_1(s)) - G(s, x_2(s))) ds \right| \\ &\leq \frac{1}{a(t)} \int_0^t a(s)L\|x_1 - x_2\|_\infty ds \\ &\leq tL\|x_1 - x_2\|_\infty < \kappa\|(x_1, y_1) - (x_2, y_2)\|_X. \end{aligned}$$

Hence, since  $T$  is a contraction, there is a unique fixed point of  $T$ , thus concluding the proof. □

Let us now face the problem of the continuous dependence on initial data. We recall that all solutions of (30) can be extended to the whole interval  $[0, 1]$  thanks to the linear growth condition (29).

**Theorem 28.** *For every  $x_0 \in \mathbb{R}$  and every  $\varepsilon > 0$  there exists  $\delta > 0$  such that, if  $\hat{x}_0 \in \mathbb{R}$  satisfies  $|x_0 - \hat{x}_0| < \delta$ , then the solution  $(x, y)$  of (30) and the solution  $(\hat{x}, \hat{y})$  of*

$$\begin{cases} x' = F(t, y), & (a(t)y)' = a(t)G(t, x), \\ x(0) = \hat{x}_0, & y(0) = 0 \end{cases}$$

satisfy

$$|x(t) - \hat{x}(t)| < \varepsilon, \quad |y(t) - \hat{y}(t)| < \varepsilon,$$

for every  $t \in [0, \rho_0]$ .

*Proof.* Fix  $x_0 \in \mathbb{R}$ . We first prove that there exists  $M > 0$  such that

$$\text{if } |x_0 - \hat{x}_0| < 1, \text{ then } |\hat{x}(t)| \leq M \text{ and } |\hat{y}(t)| \leq M, \text{ for every } t \in [0, \rho_0]. \quad (31)$$

We can compute, recalling (29), both

$$\begin{aligned} |\hat{x}(t)| &\leq |\hat{x}_0| + \int_0^t |F(s, \hat{y}(s))| ds \leq 1 + |x_0| + \int_0^t A(1 + |\hat{y}(s)|) ds \\ &\leq 1 + |x_0| + A\rho_0 + A \int_0^t |\hat{y}(s)| ds, \end{aligned}$$

for every  $t \in [0, \rho_0]$ , and, recalling assumption (A3),

$$\begin{aligned} |\hat{y}(t)| &\leq \frac{1}{a(t)} \int_0^t a(s)|G(s, \hat{x}(s))| ds \leq A\psi(t) + A\frac{1}{a(t)} \int_0^t a(s)|\hat{x}(s)| ds \\ &\leq A\rho_0 + A \int_0^t |\hat{x}(s)| ds, \end{aligned}$$

for every  $t \in [0, \rho_0]$ . Hence, setting  $z(t) = \max\{|\hat{x}(t)|, |\hat{y}(t)|\}$ , we have

$$z(t) \leq (1 + |x_0| + A\rho_0) + A \int_0^t z(s) ds,$$

so that, by Gronwall Lemma, we deduce that

$$z(t) \leq M := (1 + |x_0| + A\rho_0)e^{A\rho_0}.$$

Hence, (31) holds. Therefore, we can consider a Lipschitz constant  $L > 0$  such that, for every  $s \in [0, \rho_0]$ ,

$$\begin{aligned} |y_1| \leq M \text{ and } |y_2| \leq M &\Rightarrow |F(s, y_1) - F(s, y_2)| \leq L|y_1 - y_2|, \\ |x_1| \leq M \text{ and } |x_2| \leq M &\Rightarrow |G(s, x_1) - G(s, x_2)| \leq L|x_1 - x_2|. \end{aligned}$$

Then, we can compute, for every  $t \in [0, \rho_0]$ ,

$$\begin{aligned} |x(t) - \hat{x}(t)| &\leq |x_0 - \hat{x}_0| + \int_0^t L|y(s) - \hat{y}(s)|ds, \\ |y(t) - \hat{y}(t)| &\leq \frac{1}{a(t)} \int_0^t a(s)L|x(s) - \hat{x}(s)|ds \leq \int_0^t L|x(s) - \hat{x}(s)|ds, \end{aligned}$$

so that, defining  $z(t) = \max\{|x(t) - \hat{x}(t)|, |y(t) - \hat{y}(t)|\}$ , we find

$$z(t) \leq |x_0 - \hat{x}_0| + L \int_0^t z(s)ds,$$

and therefore

$$z(t) \leq |x_0 - \hat{x}_0|e^{L\rho_0} \leq \delta e^{L\rho_0}, \quad \text{for every } t \in [0, \rho_0].$$

Then, setting  $\delta < \min\{1, \varepsilon e^{-L\rho_0}\}$ , we conclude the proof. □

We have proved the continuous dependence on initial data for (30) in a right neighborhood of 0. Since the functions  $F$  and  $G$  satisfy (29), we can easily recover this property in the whole interval  $[0, 1]$ .

We now consider the situation treated in Sect. 3.2. To prove the continuity of the flow  $\Phi$  we need to state the analogues of Theorems 27 and 28 for the final value problems

$$\begin{cases} x' = F(t, y), & (a(t)y)' = a(t)G(t, x), \\ x(1) = x_0, & y(1) = 0, \end{cases}$$

where  $x_0 \in \mathbb{R}$ . Their proofs can be provided by the change of variable  $t \mapsto 1 - t$ .

## References

- [1] Bereanu, C., Jebelean, P., Mawhin, J.: Radial solutions for Neumann problems involving mean curvature operators in Euclidean and Minkowski spaces. *Math. Nachr.* **283**(3), 379–391 (2010)
- [2] Boscaggin, A., Colasuonno, F., Noris, B.: Multiple positive solutions for a class of  $p$ -Laplacian Neumann problems without growth conditions. *ESAIM: COCV* **24**(4), 1625–1644 (2018)
- [3] Cortázar, C., Dolbeault, J., García-Huidobro, M., Manásevich, R.: Existence of sign changing solutions for an equation with a weighted  $p$ -Laplace operator. *Nonlinear Anal.* **110**, 1–22 (2014)
- [4] De Coster, C., Habets, P.: *Two-point Boundary Value Problems: Lower and Upper Solutions*. Elsevier, Amsterdam (2006)
- [5] Fonda, A., Klun, G., Obersnel, F., Sfecci, A.: On the Dirichlet problem associated with bounded perturbations of positively- $(p, q)$ -homogeneous Hamiltonian systems. *J. Fixed Point Theory Appl.* **24**, 4 (2022)

- [6] Fonda, A., Klun, G., Sfecci, A.: Well-ordered and non-well-ordered lower and upper solutions for periodic planar systems. *Adv. Nonlinear Stud.* **21**, 397–419 (2021)
- [7] Fonda, A., Sfecci, A., Toader, R.: Two-point boundary value problems for planar systems: a lower and upper solutions approach. *J. Differ. Equ.* **308**, 507–544 (2022)
- [8] Fonda, A., Toader, R.: A dynamical approach to lower and upper solutions for planar systems. *Discrete Contin. Dyn. Syst.* **41**, 3683–3708 (2021)
- [9] Franchi, B., Lanconelli, E., Serrin, J.: Existence and uniqueness of nonnegative solutions of quasilinear equations in  $\mathbb{R}^n$ . *Adv. Math.* **118**(2), 177–243 (1996)
- [10] Nagumo, M.: Über die Differentialgleichung  $y'' = f(t, y, y')$ . *Proc. Phys. Math. Soc. Jpn.* **19**, 861–866 (1937)
- [11] Picard, E.: Sur l'application des méthodes d'approximations successives à l'étude de certaines équations différentielles ordinaires. *J. Math. Pures Appl.* **9**, 217–271 (1893)
- [12] Reichel, W., Walter, W.: Radial solutions of equations and inequalities involving the  $p$ -Laplacian. *J. Inequal. Appl.* **1**(1), 47–71 (1997)
- [13] Scorza Dragoni, G.: Il problema dei valori ai limiti studiato in grande per le equazioni differenziali del secondo ordine. *Math. Ann.* **105**, 133–143 (1931)

Alessandro Fonda, Natnael Gezahegn Mamo, Franco Obersnel and Andrea Sfecci  
 Dipartimento di Matematica e Geoscienze  
 Università degli Studi di Trieste  
 P.le Europa 1  
 34127 Trieste  
 Italy  
 e-mail: [a.fonda@units.it](mailto:a.fonda@units.it)

Natnael Gezahegn Mamo  
 e-mail: [natnaelgezahegn.mamo@phd.units.it](mailto:natnaelgezahegn.mamo@phd.units.it)

Franco Obersnel  
 e-mail: [obersnel@units.it](mailto:obersnel@units.it)

Andrea Sfecci  
 e-mail: [asfecci@units.it](mailto:asfecci@units.it)

Natnael Gezahegn Mamo  
 Department of Mathematics  
 Addis Ababa University  
 Adwa Street  
 1176 Addis Ababa  
 Ethiopia

Received: October 12, 2022.

Revised: January 19, 2023.

Accepted: February 16, 2023.